



EFFECT OF PESTICIDES ON SOIL ECOSYSTEM SERVICES AND PROCESSES

MONICA JAT*, PYNHUNLIN NOLA KHARKRANG DOHLING, AMIT AHUJA^{1*} AND JAGMOHAN SINGH²

Division of Entomology; ¹Division of Nematology; ²Division of Plant Pathology,
ICAR-Indian Agricultural Research Institute, New Delhi 110012, India

*Email: monicajat174@gmail.com & aahuja165@gmail.com (corresponding authors)

ABSTRACT

The livelihood of the entire population and the existence of farming communities is solely dependent on agriculture. Fertility of the soil is governed by its texture and diversified fauna present in the soil medium. Various agricultural inputs are applied with the eventual aim to achieve the highest productivity whereas hazardous effects on soil microorganisms are often ignored. Pesticides are noxious chemicals used to mitigate crop losses incurred by nematodes, insects, weeds, mites, and pathogens. Fungicides modify the fungi communities accountable for the nutrient recycling in the soil, especially copper-based fungicides which cause harmful effects on earthworms. The continuous use of herbicides indirectly reduces the associated arthropod communities. Insecticides cause toxic effects on most of the lower soil invertebrates and cause negative impact on their growth and development and alter the trophic structure in their habitats. Buildup of synthetic pesticides in the soil cause many sublethal effects on the existing natural fauna and affects their survival. However, pesticides help in diminishing the quantitative losses caused by pests and enhance the agricultural productivity, but consequently the cost of enhanced productivity is compensated by the Earth's ecosystems. This review provides insight into the effects of pesticides on the soil health.

Key words: Ecosystem services, communities, insects, nematodes, weeds, mites, pathogens, pesticides, insecticides, herbicides, fungicides, nematicides

Pesticides are used to safeguard the crops against various threatening pests, which include insects, nematodes, mites and rodent etc. In India, the losses due to pests stand for US\$ 42.66 million in monetary terms (Sushil, 2016). Pesticides may be organic or inorganic compounds and categorized as per target pests. Pesticides may be an insecticide, herbicides, rodenticides, molluscicides, and nematicides etc. are the essential evil as they can manage the pest's problem rapidly in the field compared to all other available management techniques. But on the other hand, these pesticides are accountable for the weakening of the environment and human health; have a threat to essential fauna and flora existing in aquatic and terrestrial ecosystems. Any release of new pesticides goes through various developmental processes of risk analysis on essential micro-fauna and micro-flora of soil and water. Microbial densities and their activities determine the physical properties of soil (Abd-Alla and Omar, 1993), which possess an unavoidable threat from the pesticide residues in soil or in groundwater. The long-term persistence of pesticides residues impacts negative effects on the soil function and properties like rhizodeposition, soil organic carbon, pH, and moisture etc. The microbial populations or decomposers are

responsible for the recycling of net primary production (Hussain et al., 2009) and participate in various soil biological processes (Cycon and Piotrowska-Seget, 2007). But microbial population stability is disturbed by erratic and excess uses of pesticides. The arbuscular mycorrhiza fungi are responsible for the better growth and development of plants but reported to be affected by pesticide's toxicity, which suppresses the growth of mycelia (Sreenivasa and Bagyaraj, 1989) and systemic pesticides damage AM fungi more than the non-systemic fungi (Menge, 1982). It has been reported that nodulation in legume crops by *Rhizobium* and *Bradyrhizobium* bacteria is suppressed by the over uses of pesticides (Abd-Alla et al., 2000).

Overview of pesticide use in agriculture in India

In India, the insecticide and fungicide market is dominant over herbicides and rodenticides (Fig. 1). The share of insecticides reduced from 70% in 2003-04 to 39% in 2016-17. Indian traders generally procure technical ingredients of the pesticides formulations, which is safeguarded by patents and the readymade formulations are exported to many countries via established trade channels. Average consumption of pesticide in India is 0.6 kg/ha as per CIB&RC, which




Fig. 1. Usages of pesticides in agriculture (Subash et al., 2017)

is very low in comparison to developed countries. In the category of pesticides, insecticides shares are highest among all pesticides. The overall uses of pesticides reported highest in Maharashtra, followed by Madhya Pradesh, Punjab and Haryana. Consumption of pesticides is highest in Punjab which is reported to be approximately 0.74 kg/ha (Fig. 2).




Fig. 2. Chronological changes in total consumption of pesticides in India (Subash et al., 2017)

Ecosystem services and disservices to agriculture

The physiochemical properties of the soil are governed by the farming practices adopted by the farmers during crop cultivation. Many live organisms render their services to improve soil structure and texture. The earthworm enhances the soil structure through soil burrows which creates the pores and by producing casts and increase fertility through partial digestion (Edwards, 2004). Even many insects render their ecosystem services by facilitating pollination, by decomposition of waste material, and recycling of nutrition etc. (Losey and Vaughan, 2006). The genetic diversity also plays an important role in rendering raw material for natural selection, which helps in evolutionary adaptations (Hawtin, 2000). The indiscriminate and erratic use of synthetic pesticides is responsible for pest outbreaks, resurgence, sudden enhancement in fecundity, and resistance in pest population against

pesticides (Thomas, 1999). The ecosystem service and dis-services are depicted in Fig. 3.

The fate of pesticides in the environment

In the environment, the assessment and evaluation of fate of pesticides are essential to determine, as they play an important role in suppressing pest population, simultaneously cause harm to the beneficial microbial population and intoxicate the aquatic and terrestrial resources (Aktar et al., 2009). An ideal pesticide should only be lethal to the target pest, easily biodegradable by the microbes, and should exhibit negligible leaching in the geosphere, but this seems impossible in maximum cases. The fate of pesticides is governed by number of factors such as absorption and uptake by plants, soil adsorption, soil leaching, water runoff, volatilization and wind drift (Aktar et al., 2009). The chance of pesticide movement to off-site is dependent on the formulations, dose rates, chemical properties, modes of application, soil properties, frequency of rainfall or irrigation, and depth to groundwater etc. (Chakrabarti et al., 2006; Aldrich, 2011). The mechanism of retaining pesticides on the soil particle surfaces is adsorption, dependent on the chemical properties of pesticide and soil particulate matters (Aktar et al., 2009). Anionic and neutral pesticides are mostly mobile in soils whereas cationic pesticides exhibit strong adhesion to soil particles (Aldrich, 2011).

Factors affecting toxicity

The application dosage is the most important factor which determines the pesticide's residue in the environment. Lower dosages of pesticides are mostly non-toxic or cause slighter negative effects on the microbial fauna in soil but as the application dosage increase, concentration of residues and the toxicity level also increases. Wang et al. (2008) reported that the excessive use of pesticides such as methamidophos adversely affected the microbial mass and composition, genetic diversity and also the catabolic activities those were estimated by several culturing technique as phospholipid derived fatty acid (PLFA), community-level catabolic profiles (CLCP), amplified ribosomal DNA restriction analysis (ARDRA). The herbicides like glyphosate reduce the population of actinomycetes in humus soil (chernozem), but in wetland soil (gleysol), it enhances their population (Sumalan et al., 2010). The toxicity of pesticides is also impacted by other factors includes various kinds of application methodology like pre-seed or repetitive treatments or by addition of organic manures and on the specific age of the




Fig. 3. Ecosystem services and dis-services to agriculture (Zhang et al., 2007)

crop plant (Kalyanasundaram and Kavitha, 2012). It has been reported that the functional biodiversity of bacteria in the soil is reduced by the pre and after seed treatments.

Factors affecting the water pollution

The rate and magnitude of pesticide flow is influenced by both pesticide properties as well as soil properties. The mobility of hydrophobic pesticides and hazard of their discharge into groundwater is addressed by the weak force between pesticide particles and the soil matrix (Wauchope et al., 2001). The hydrogeological conditions of the soil, land use pattern, type of pollutants are responsible for groundwater pollution (Woof et al., 1999). Groundwater vulnerability is a term used to represent the natural ground characteristics with which groundwater may be contaminated by human activities. (US EPA) United States Environment Protection Agency and (NRC) National Research Council (1993) used an idea of groundwater susceptibility along with aquifer sensitivity which does not include characteristics of pesticides and land use or field practices and US EPA developed a DRASTIC system for evaluation of groundwater pollution potential (Aller et al., 1987).

Effect on soil ecosystem services

Pesticides manipulate the nodulation and biological nitrogen fixation in legumes by affecting the infection of nodular bacteria in root fibers of the plant. Herbicide affects the nodulation by adverse effects on rhizobia and plant growth (Zawoznik and tomoro, 2005). Soil profiles possess mobile enzymes and static extracellular enzymes which are an indicator of biological stability, fertility and quality and change in biological status due to pollution (Mayanglambam et al., 2005). Nitrogenase is an enzyme required by the organisms to fix atmospheric nitrogen. Dehydrogenase enzymes occur inside the living microbial cells which are responsible for microbial respiratory processes. The breakdown and alteration of organic matter is carried out by various enzymes as β - Glucosidase, cellulase, and invertase. The urea is hydrolysed into CO_2 and NH_3 , a key component of N- cycle is catalyzed by urease enzymes. In Table 1 and 2, a brief summary of effects of various pesticides on biochemical reactions and soil enzymes is listed, respectively.

Microbes

Soil organisms mainly consist of soil microbes like

Table 1. Effects of various pesticides on biochemical reactions in soil

Pesticide	Biochemical reaction	Comments	References
BHC, fenvelerate	C- mineralization	Increased	(Murthy et al., 1991)
Dithiocarbamate	Nitrification	Fungicides inhibited ammonium oxidation	(Hanson et al., 1991)
Chlorinated hydrocarbons	Methanogenesis	Inhibited	(Krumback and Conrad, 1991)
Fenpropimorph	Denitrification	Inhibited	(Svensson and Leonardsson, 1992)
Fungicides	Nitrification	Inhibited nitrification of urea	(Martens and Bremner, 1993)
Cyfluthrin, imidacloprid	Nitrification, S-oxidation, denitrification	Stimulated S- oxidation and inhibited nitrification and denitrification	(Tu, 1995)
Dinoterb	N-mineralization	Enhanced	(Engelen et al., 1998)
Terbutryn, simazine, premetryn, benzoate	N-fixation	Decrease N content and nodulation	(Singh and Wright, 1999)
Butaclor	N-fixation	Initially Increased and later decrease	(Min et al., 2001)
Bensulfuron methyl	N-mineralization	Decreased	(El- Ghamry et al., 2001)
Captan, Benomyl, Chlorothalonil, anilazine	N- mineralization	Increased	(Chen and Edwards, 2001)
DDT, 2-4 D, 2-4 T	N- fixation	Inhibited nodulation	(Fox et al., 2001)
Metalaxyl, Mefonoxam	Ammonification, Nitrification	Increased	(Monkiedje et al., 2002)
Mancozeb, prosulfuron, chlorothalonil	Nitrification	Inhibited	(Arora et al., 2003)
Chlorpyrifos, Quinalphos	Ammonification	Inhibited	(Menon et al., 2004)
Carbendazim, Thiram, Imazethpyr	N-fixation	Inhibit nodulation	(Niewiadomska, 2004)
Pentachlorophenol	Nitrification	Reduced	(Colores and Schmidt, 2005)
Acetamiprid	Respiration	Reduced	(Yao et al., 2006)
Glyphosate, Imidacloprid and hexaconazole	N-fixation	Toxic to <i>Bradyrhizobium</i> sp. strain MRM6	(Ahemad and Khan, 2011)
Fenamiphos	Nitrification	Inhibited	(Caceres et al., 2014)

bacteria and fungi, protozoa, nematodes, earthworms and mites etc. Soil organisms have a great influence on nutrient availability by governing number of factors such as mineralization, nutrient immobilization, carbon content in soil, crop residues decomposition and bioturbation. The microfauna has a greater group of enzymes that plays important role in the chemical disintegration of organic matter (Paul and Clark, 1996). Soil microbial biomass which majorly comprises of soil inhabiting bacteria and fungi those are responsible for mineralization and immobilization of soil nutrients. Soil microbial biomass is considered as the source of crucial primary nutrients (C, N, P, S) those are released through their grazing activities of soil fauna. Most prominent method to measure the outcome of pesticides on the soil organism is to determine the change in biological activities i.e. by soil respiration, enzymatic activity or microbial biomass. Effects of various pesticides on the functioning of soil microbial fauna are listed in Table 3.

Collembola

Collembolans are free-living organisms, present in

all ecosystem mostly prefer humid conditions. They are responsible for the fragmentation of organic matter and indirectly involved in the decomposition of O.M. Usually, they prefer to consume fungal hyphae and spores, but sometimes also bacteria or plant material. In the terrestrial ecosystem, three invertebrate species are well-known for ecotoxicological assays are *Eisenia fetida* (Lumbricidae), *Folsomia candida* (Collembola), and *Enchytraeus albidus* (Enchytraeidae), as all species has a short life cycle as well as parthenogenetic reproduction. All these qualities make it appropriate for assays (Jansch et al., 2006). Collembolans has less contribution in soil respiration processes and biomass (Coleman et al., 2004) but affecting disintegration of O.M and nutrient cycling and has major influence on microbial ecology and soil fertility (Culik and Zeppelini, 2003). Alves et al. (2014) investigated the effects of seed dressing pesticides on survival and reproduction of *Folsomia candida*, by ecotoxicological tests. Five seed dressing pesticides as imidacloprid, fipronil, thiamethoxam, captan and carboxin+ thiram. Imidacloprid and fipronil were lethal even at lower concentrations which prove as the most toxic pesticides,

Table 2. Effects of pesticides on soil enzymes

Pesticide	Soil enzymes	Comments	References
Captan, thiram	Nitrogenase	Activity of <i>Azospirillum brasilense</i> was inhibited	Gallori et al., 1991
Fenvalerate, cuprosan	Nitrogenase	Inhibited	Omar and Abd-alla, 1992
Profenophos	Nitrate reductase	Inhibited	Abdel-Malik et al., 1994
Captan	Nitrogenase	Inhibited	Di Ciocco and Caceres, 1997
Terbutryn, simazine, prometryn	Nitrogenase	Reduced	Singh and Wright, 1999
Glyphosate	Dehydrogenase	At first inhibited but afterwards restored	Andrea et al., 2000
Brominal, selecron	Cellulase	Inhibited	Omar and Abdel-Sater, 2001
Oxafun, Funaben, Baytan	Nitrogenase	Higher dose reduced the activity	Durska, 2004
Carbendazim, imazetapyr, thiram	Nitrogenase	Reduced	Niewiadomska, 2004
Isoproturon	Dehydrogenase	Decrease	Nowak et al., 2004
Isoproturon	Urease	Increase	Nowak et al., 2004
Metalaxyl	Urease, Phosphatase	Decrease and first increase and then decreased respectively	Sukul, 2006
Methabenzthiazuron, terbutryn	Nitrogenase	Adversely affect enzymatic activity and reduce nodule number.	Khan et al., 2006
Atrazine, northrin	Dehydrogenase	At low concentration, activity stimulated while at higher concentration it reduces	Nweke et al., 2007
Azoxystrobin, tebuconazole, chlorothalonil	Dehydrogenase	In the low OM soil, activity reduces but not in high OM soil	Bending et al., 2007
Validamycin	Phosphatase and urease	Validamycin affect soil enzyme but recovered soon	Qian et al., 2007
Fenamiphos	Dehydrogenase and urease	Not toxic	Caceres et al., 2009
Endosulfan	Dehydrogenase	Increased	Kalyani et al., 2010
Diuron	Urease	No effect	Romereo et al., 2010
Propiconazole	Cellulase	Reduced from 5-40%	Ramudu et al., 2011
Thiamethoxam	Dehydrogenase, urease and phosphatase	Inhibitory effect on dehydrogenase and phosphatase	Jyot et al., 2015
Azoxystrobin	Urease, dehydrogenases, acid phosphatase, catalase and alkaline phosphatase	Dehydrogenases-resistant while alkaline phosphatase recovered in the soil, rest show inhibitory effect	Bacmaga et al., 2015
Dimethomorph	Dehydrogenases, urease, invertase and acid phosphatase	Urease- no effect Dehydrogenase- Inhibited Invertase- stimulated Acid phosphatase- no effect	Wang et al., 2016

reducing significantly the collembolan reproduction. Frampton and Brink (2007) studied direct and indirect effects of carbamate, synthetic pyrethroid and organophosphate insecticides on collembola and macroarthropods community, suggested that cypermethrin induced the abundance of collembolan, represent a classical resurgence while primicarb does not affect collembolans.

Mycorrhiza

Mycorrhiza exhibit symbiotic association between

a green plant and a fungus. The plant produces sugar by photosynthesis and supplies it to fungi, and in support fungi supply plant water, nutrients as phosphorus from soil. Many experiments revealed that legumes inoculated with both rhizobium and arbuscular mycorrhiza (AM) has an additive effect on plant growth rather than inoculated individually (Vejsadova et al., 1989). It is responsible for increase in P uptake and nitrogen fixation which provides a supplementary benefit towards the synergistic relationship (Brown and Bethlenfalvay, 1987). Abd-alla et al. (2000) has studied

Table 3. Effects of various pesticides on soil microbial fauna

Pesticides	Comments	References
Captan	Fungal hyphae length and density, reduction in microbial C and N	Hu et al., 1995
Copper	Soil microbial biomass reduced	Belotti, 1998
DDT	Have adverse effect on bacteria and algal biomass and may not have any impact on fungal count	Meghraj et al., 2000
Captan, chlorothalonil	Suppression of microbial respiration	Chen et al., 2001
Copper	Metabolic quotient was increased which indicates microbial stress. Ratio of microbial biomass to organic carbon was significantly reduced	Merrington et al., 2002
Metalaxyl, Mefenoxam	Bacteria numbers increased with high doses, but toxic towards N fixers	Monkiedje et al., 2002
Glyphosate	Significant reduction in bacteria and fungi population, but increase in actinomycetes. Microbial activity enhanced by 9–19%	Araujo et al., 2003
DDT, arsenic contamination	Significant reduction in the bacterial and fungal count, and biomass carbon was observed	Edvantro et al., 2003
Atrazine, metolachlor	Change in the community structure of actinomycetes and bacteria	Seghers et al., 2003
Arsenic	Decline in the microbial biomass, dehydrogenase activity and soil respiration	Ghosh et al., 2004
Pendimethalin	Decline in the population of nematodes and plant rhizobium in the soil	Strandberg and Scott-Fordsmann, 2004
Chlorpyrifos, dimethoate	Chlorpyrifos has, has the more lethal effect towards collembolan density	Endlweber et al., 2005
Chlorothalonil, azoxystrobin	Chlorothalonil and azoxystrobin were found toxic to the <i>Fusariumoxysporum</i> strain CS-20 (biocontrol agent)	Fravel et al., 2005
Atrazine and metolachlor	Reduction in the microbial counts	Ayansina and Oso, 2006
Diazinon, linuron and mancozeb+ dimethomorph	CFU (colony forming unit) of bacteria and fungi decreased	Cycon and Seget, 2007
Carbofuran, butachlor, diuron, Chlorotoluron, glyphosate, methamidophos, fenamiphos	Carbofuran, glyphosate, methamidophos stimulate while butachlor and femaiphos were detrimental to microbial growth	Lo, 2010
Cypermethrin and thiomethoxam	Detrimental effect on metabolic processes of soil microorganism	Filimon et al., 2015

the effects of various pesticides like Afugan, Brominal, Gramoxone, Selecron and Sumi Oil on development, nodulation and root colonization of the cowpea (*Vigna sinensis* L.), common bean (*Phaseolus vulgaris* L.) and Lupin (*Lupinus albus* L.) by AM. Application of pesticides restricts the growth of all the legumes but vary with the pesticide and plant species. Growth of the some legumes crops is harmfully exaggerated by indiscriminate and erratic use of pesticides but the extent of the effect modifies with type of plant species and class of pesticides used. In cowpea, after 20 (DAP) days of planting, nodule formation was significantly inhibited and recovered at 40 and 60 days after planting. In all legumes except cowpea, AM root colonization and spores are inhibited, but spore formation was increased in pesticide treated cowpea after 60 DAP.

Earthworms

Earthworms are considered as soil engineers as they are usually present in soil, consume organic matter like plant matter and microfauna. Earthworms

are classified in to 3 main ecophysiological categories, epigenic, endogenic and anecic. Epigenic earthworms e.g. *Eisenia fetida*, are the non burrowing types and live in the soil litter boundary and consume disintegrated O.M. Endogenic earthworms dwell in the subsoil and makes flat burrows in 10–30 cm of soil, while Anecic earthworms e.g. *Lumbricus terrestris* construct vertical burrows in search of food. Their population depends on soil physiochemical properties such as temperature, pH, salts and moisture. Mostly species are present in neutral to slightly acidic soils. If the soil is more acidic, earthworm would go in to diapause stage easily. Soil acidity is increased by the use of nitrogenous fertilizers, which prove lethal to worms. The effects of various pesticides on biology of earthworm are listed in Table 4.

CONCLUSIONS

Agriculture is the mean to serve the mankind by providing food and generate employment and the top soil is the just the medium to sustain the agriculture. The

Table 4. Effects of pesticides on biology of earthworms

Organisms	Comments	References
<i>Eisenia fetida</i>	Isoproturon affects earthworm at very high concentration	Mosleh et al., 2003
<i>Drawida willsi</i>	Butachlor is reported as lethal to earthworm <i>Drawida willsi</i> by have harmful effect on growth and development, sexual maturity and cocoon formation	Panda and Sahu, 2004
Earthworm sp.	Growth rate and borrowing activity of earthworms is reduced with copper which leads to increase in soil bulk densities	Eijsackers et al., 2005
<i>Eisenia fetida</i>	Malathion decreased the spermatocytic viability in spermatheca and modifying the DNA structure of spermatogonia	Navarro and Obregon, 2005
<i>Apporectodea nocturna</i> and <i>Allolobophora lcterica</i>	Imidacloprid, reduced burrowing activity in both earthworm species	Capoweiz and Berard, 2006
<i>Eisenia andrei</i>	Malathion + primiphos methyl show antagonistic effect against <i>Eisenia andrei</i>	Stepic et al., 2013
<i>Eisenia fetida</i>	Imidacloprid has a harmful effect on <i>E. fetida</i> as it affects the antioxidant defense and digestive system	Zhang et al., 2014
<i>Eisenia fetida</i>	The toxicity of 5 pesticides (trichlorfon, dimethoate, carbendazim, tebuconazole and prochloraz) was evaluated on different lethal and sublethal end points of the earthworm	Rico et al., 2016
<i>Eisenia fetida</i>	The herbicide tribenuron-methyl and fungicide tebuconazole have exhibited antagonistic effects on soil earthworm <i>Eisenia fetida</i>	Chen et al., 2018

topsoil needs utmost care as the quality and fertility of the soil determines the yield of crop. By adaptation of modern days tactics like uses of synthetic fertilizers and pesticides, have enhanced the yield output, but the same time many problems related to health of soil-ecosystem have evolved. The directive of agricultural development is to nourish and deliver surplus nutrition to the growing population without negotiating with the health of environment. Widespread and erratic application of synthetic chemical fertilizers and pesticides and their leaching in to groundwater lead to soil and water pollution. In cultivated lands, the soil ecosystem services provided by microfauna are affected by the application of toxic and persistent chemical pesticides which ultimately changes the soil properties. Synthetic pesticides are harmful to soil ecosystem and human health as derivatives of pesticides are persistent and long lasting in soil which negatively affect the growth and development, diversity and biochemical processes of microbes. Pesticides also reduce soil fertility by which crop yield is affected. A rational use of the pesticides is important as harmful effects are observed when the applied doses are more than the recommended rates. To reduce the hazardous effects on the environment and humans, the farmers, distributors, industry, stakeholders and policymakers should educated and trained for rational use of pesticide. Soil microbial population provides a vital link in the natural food web. Thus, any alteration in the composition and soil microbial population can disturb the trophic structure and cause a risk to global food security. Well-designed trials and experiments are the need of the hour to examine the

long term consequences of the pesticides on the health of the microbial communities as well as their ecotoxicological effects in the soil environment.

REFERENCES

- Abd-Alla M H, Omar S A. 1993. Herbicides effects on nodulation, growth and nitrogen yield of faba bean induced by indigenous *Rhizobium leguminosarium*. Zentralbl Mikrobiol 148: 593-597.
- Abd-alla M H, Omar S, Karanxha S. 2000. The impact of pesticide on arbuscular mycorrhizal and nitrogen fixing symbioses in legumes. Applied Soil Ecology 14(3): 191-200.
- Abdel-Mallek A Y, Moharram A M, Abdel-Kader M I, Omar S A. 1994. Effect of soil treatment with the organophosphorus insecticide profenofos on the fungal flora and some microbial activities. Microbiology Research 149: 167-171.
- Ahemad M, Khan M S. 2011. Effect of pesticides on plant growth promoting traits of greengram-symbiont, *Bradyrhizobium* sp. strain MRM6. Bulletin of Environmental Contamination and Toxicology 86: 384-388.
- Ahemad M, Khan M S. 2013. Pesticides as antagonists of rhizobia and the legume-rhizobium symbiosis: a paradigmatic and mechanistic outlook. Biochemistry and Molecular Biology 1(4): 63-75.
- Aktar M W, Sengupta D, Chowdhury A. 2009. Impact of pesticides use in agriculture: their benefits and hazards. Interdisciplinary Toxicology 2: 1-12.
- Aldrich S. 2011. Material Safety Data Sheets. Sigma-Aldrich, Oakville, Ontario.
- Aller L, Bennett T, Lehr J H, Petty R J, Hackett G. 1987. DRASTIC: A standardised system for evaluating groundwater pollution potential using hydrogeologic settings. US Environment Protection Agency, Ada, Oklahoma EPA/60012-87/035.
- Alves P L R, Cardoso E J B N, Alexandre M M, Sousa J P, Pasini A. 2014. Seed dressing pesticides on springtails in two ecotoxicological laboratory tests. Ecotoxicology and Environmental Safety 105: 65-71.
- Andrea M M, Peres T B, Luchini L C, Pettinelli A Jr. 2000. Impact of

- long-term pesticide application on some soil biological parameters. *Journal of Environment Science Health Biology* 35: 297-307.
- Araujo A S F, Monteiro R T R, Abarkeli R B. 2003. Effect of glyphosate on the microbial activity of two Brazilian soils. *Chemosphere* 52: 799-804.
- Arora A, Singh B, Sud D, Srivastava T, Arora C L. 2003. Inhibition of nitrification in soil by metal diethylthiocarbamates. *Journal of Environment Science* 15: 607-610.
- Ayansina A D V, Oso B A. 2006. Effect of two commonly used herbicides on soil microflora at two different concentrations. *African Journal of Biotechnology* 5(2): 129-132.
- Bacmaga M, Kucharski J, Wyszowska J. 2015. Microbial and enzymatic activity of soil contaminated with azoxystrobin. *Environment Monitoring Assess* 187: 615.
- Belotti E. 1998. Assessment of a soil quality criterion by means of a field survey. *Applied Soil Ecology* 10: 51-63.
- Bending G D, Rodr'iguez-Cruz M S, Lincoln S D. 2007. Fungicide impacts on microbial communities in soils with contrasting management histories. *Chemosphere* 69: 82-88.
- Brown M S, Bethlenfalvai G J. 1987. Glycine-Glomus-Rhizobium symbiosis. VI. Photosynthesis in nodulated mycorrhizal, or N- and P-fertilized soybean plants. *Plant Physiology* 85: 120-123.
- Caceres T P, He W, Megharaj M, Naidu R. 2009. Effect of insecticide fenamiphos on soil microbial activities in Australian and Ecuadorean soils. *Journal of Environmental Science and Health Biology* 44: 13-17.
- Capowiez Y, Berard A. 2006. Assessment of the effects of imidacloprid on the behavior of two earthworm species (*Aporrectodea nocturna* and *Allolobopora*) using 2D terraria. *Ecotoxicology and Environmental Safety* 64: 198-206.
- Chakrabarti K, Chakraborty A, Chowdhury A. 2006. Degradation and effects of pesticides on soil microbiological parameters. *International Journal of Agricultural Research* 1: 240-258.
- Chen J, Saleem M, Wang C, Liang W, Zhang Q. 2018. Individual and combined effects of herbicide tribenuron-methyl and fungicide tebuconazole on soil earthworm *Eisenia fetida*. *Scientific reports* 8: 29-67.
- Chen S K, Edwards C A. 2001. A microcosm approach to assess the effects of fungicides on soil ecological processes and plant growth: Comparison of two soil types. *Soil Biology and Biochemistry* 33: 1981-1991.
- Coleman D C, Crossley D A, Hendrix P F. 2004. *Fundamentals of soil ecology* (second edition).
- Colores G M, Schmidt S K. 2005. Recovery of microbially mediated processes in soil augmented with a pentachlorophenol-mineralizing bacterium. *Environment Toxicology Chemistry* 24: 1912-1917.
- Culik M P, Zeppelini Filho D. 2003. Diversity and distribution of Collembola (Arthropoda: Hexapoda) of Brazil. *Biodiversity and Conservation* 12: 1119-1143.
- Cycon M, Piotrowska-Seget Z. 2007. Effect of selected pesticides on soil microflora involved in organic matter and nitrogen transformations: pot experiment. *Polish Journal of Ecology* 55(2): 2017-220.
- Di Ciocco C A, Caceres E R. 1997. Effect of fungicide captan on *Azospirillum brasilense* Cd in pure culture and associated with *Setaria italica*. *Review of Argentina Microbiology* 29: 152-156.
- Durska G. 2004. Fungicide effect on nitrogenase activity in methylophilic bacteria. *Polish Journal of Microbiology* 53: 155-158.
- Edvantoro B B, Naidu R, Megharaj M, Singleton I. 2003. Changes in microbial properties associated with long-term arsenic and DDT contaminated soils at disused cattle dip sites. *Ecotoxicology and Environmental Safety* 55: 344-351.
- Edwards C. 2004. *Earthworm ecology*. CRC Press, Boca Raton FL 441.
- Eijsackers H, Benke P, Maboeta M, Louw J P E, Reinecke A J. 2005. The implications of copper fungicide usage in vineyards for earthworm activity and resulting sustainable soil quality. *Ecotoxicology and Environmental Safety* 62: 99-111.
- El-Ghamry A M, Huang C Y, Xu J M. 2001. Combined effects of two sulfonylurea herbicides on soil microbial biomass and N-mineralization. *Journal of Environment Science* 13, 311-317.
- Endlweber K, Schadler M, Scheu S. 2005. Effects of foliar and soil insecticide applications on the collembolan community of an early set-aside arable field. *Applied Soil Ecology* 31: 136-146.
- Engelen B, Meinken K, Von Wintzingerode F, Heuer H, Malkomes H P, Backhaus H. 1998. Monitoring impacts of a pesticide treatment on bacterial soil communities by metabolic and genetic fingerprinting in addition to conventional testing procedures. *Applied Environment Microbiology* 64: 2814-2821.
- Estevez M A, Periago E P, Martinez-Carballo E, Simal-Gandara J, Mejuto J C, Garcia-río L. 2007. *Agriculture, Ecosystem and Environment* 123 (2008): 247-260.
- FAOSTAT. 2017. *Pesticides*. Food and Agriculture Organization, Rome.
- Filimon M N, Voia S O, Popescu R, Dumitrescu G, Ciochina C P, Mituletu M, Vlad D C. 2015. The effect of some insecticides on soil microorganisms based on enzymatic and bacteriological analyses. *Romanian Biotechnological Letters* 20(3): 10439-47.
- Fox J E, Starcevic M, Kow K Y, Burow M E, McLachlan J A. 2001. Nitrogen fixation: Endocrine disrupters and flavonoid signaling. *Nature* 413: 128-129.
- Frampton G K, van den Brink P J. 2007. Influence of cropping on the species composition of epigeic Collembola in arable fields. *Pedobiologia* 46: 328-337.
- Fravel D R, Deahl K L, Stommel J R. 2005. Compatibility of the biocontrol fungus *Fusarium oxysporum* strain CS-20 with selected fungicides. *Biological Control* 34: 165-169.
- Gallori E, Casalone E, Colella C M, Daly S, Polsinelli M. 1991. 1,8-Naphthalic anhydride antidote enhances the toxic effects of captan and thiram fungicides on *Azospirillum brasilense* cells. *Res. Microbiology* 142: 1005-1012.
- Ghosh A K, Bhattacharyya P, Pal R. 2004. Effect of arsenic contamination on microbial biomass and its activities in arsenic contaminated soils of Gangetic West Bengal, India. *Environment International* 30: 491-499.
- Groffman P M, Driscoll C T, Likens G E, Fahey T J, Holmes R T, Eagar C, Aber J D. 2004. No gloom of night: a new conceptual model for the Hubbard Brook Ecosystem Study. *Bioscience* 54: 139-148.
- Hanson G B, Klemedtsson L, Stenstrom J, Torstenson L. 1991. Testing the influence of chemicals on soil autotrophic ammonium oxidation. *Environment Toxicology and Water Quality* 6: 351-360.
- Hawtin G C. 2000. Genetic diversity and food security. *The UNESCO Courier* (May 2000). pp. 27-29.
- Hu S, Coleman D C, Hendrix P F, Beare M H. 1995. Biotic manipulation effects on soil carbohydrates and microbial biomass in a cultivated soil. *Soil Biology and Biochemistry* 27: 1127-1135.
- Hussain S, Siddique T, Saleem M, Arshad M, Khalid A. 2009. Impact of pesticides on soil microbial diversity, enzymes, and biochemical reactions. *Advances in Agronomy* 102, ISSN 0065-2113.
- Jansch S, Frampton G K, Rombke J, van den Brink P J, Scott Fordsmand J J. 2006. Effects of pesticides on soil invertebrates in model

- ecosystem and field studies: a review and comparison with laboratory toxicity data. *Environment Toxicology Chemistry* 25: 2490-2501.
- Jyot G, Mandal K, Singh B. 2015. Effect of dehydrogenase, phosphatase and urease activity in cotton soil after applying thiamethoxam as seed treatment. *Environment Monitoring Assessment* 187: 298.
- Kalyanasundaram D, Kavitha S. 2012. Effect of butachlor on the microbial population of direct sown rice. *World Academy of Science and Engineering Technology* 69: 853-855.
- Kalyani S S, Sharma J, Dureja P, Singh S, Lata. 2010. Influence of endosulfan on microbial biomass and soil enzymatic activities of a tropical Alfisol. *Bulletin of Environment Contamination Toxicology* 84: 351-356.
- Khan M S, Zaidi A, Rizvi P Q. 2006. Biotoxic effects of herbicides on growth, nodulation, nitrogenase activity, and seed production in chickpeas. *Communication of Soil Science Plant Analysis* 37: 1783-1793.
- Krumback M, Conrad R. 1991. Metabolism of position labelled glucose in anoxic methanogenic paddy soil and lake sediment. *FEMS Microbiology Ecology* 85: 247-256.
- Lo C C. 2010. Effect of pesticides on soil microbial community. *Journal of Environmental Science and Health Part B Pesticides Food Contaminants and Agricultural Wastes* 45(5): 348-359.
- Losey J E, Vaughan M. 2006. The economic value of ecological services provided by insects. *Bioscience* 56 (4): 331-323.
- Martens D A, Bremner J M. 1993. Influence of herbicides on transformations of urea nitrogen in soil. *Journal of Environment Science Health* 28: 377-395.
- Mayanglambam T, Vig K, Singh D K. 2005. Quinalphos persistence and leaching under field conditions and effects of residues on dehydrogenase and alkaline phosphomonoesterases activities in soil. *Bulletin of Environment Contamination Toxicology* 75: 1067-1076.
- Megharaj M, Kantachote D, Singleton I, Naidu R. 2000. Effects of long-term contamination of DDT on soil microflora with special reference to soil algae and algal transformation of DDT. *Environmental Pollution* 109: 35-42.
- Menge J A. 1982. Effect of soil fumigants and fungicides on vesicular–arbuscular fungi. *Phytopathology* 72: 1125-1132.
- Menon P, Gopal M, Parsad R. 2004. Influence of two insecticides, chlorpyrifos and quinalphos, on arginine ammonification and mineralizable nitrogen in two tropical soil types. *Journal of Agricultural Food Chemistry* 52: 7370-7376.
- Merrington G, Rogers S L, Zwieten L V. 2002. The potential impact of long-term copper fungicide usage on soil microbial biomass and microbial activity in an avocado orchard. *Australian Journal of Soil Research* 40: 749-759.
- Min H, Ye Y F, Chen Z Y, Wu W X, Yufeng D. 2001. Effects of butachlor on microbial populations and enzymes activities in paddy soil. *Journal of Environment Science Health* 36: 581-595.
- Monkiedje A, Spiteller M. 2002. Effects of the phenylamide fungicides, mefenoxam and metalaxyl, on the biological properties of sandy loam and sandy clay soils. *Biology of Fertile Soils* 35: 393-398.
- Mosleh Y Y, Paris-Palacios S, Couderchet M, Vernet G. 2003. Effects of the herbicide isoproturon on survival, growth rate, and protein content of mature earthworms (*Lumbricus terrestris* L.) and its fate in the soil. *Applied Soil Ecology* 23: 69-77.
- Murthy N B K, Kale S P, Raghu K. 1991. Mineralization of 14C-labelled rice straw in aerobic and anaerobic clay soils as influenced by insecticide treatment. *Soil Biology and Biochemistry* 23: 857-859.
- National Research Council 1993. Ground water vulnerability assessment: Contamination potential under conditions of uncertainty. National Academy Press, Washington, DC.
- Navarro O E, Obregon E B. 2005. Effect of malathion on the male reproductive organs of earthworms, *Eisenia foetida*. *Asian Journal of Andrology* 7(1): 97-101.
- Niewiadomska A. 2004. Effect of carbendazim, imazetapyr and thiram on nitrogenase activity, the number of microorganisms in soil and yield of red clover (*Trifolium pretense* L). *Pollution Journal of Environment Studies* 13: 403-410.
- Nowak A, Nowak J, Klodka D, Pryzbulewska K, Telesinski A, Szopa E. 2004. Changes in the microflora and biological activity of the soil during the degradation of isoproturon. *Journal of Plant Disease Protection* 19: 1003-1016.
- Nweke C O, Ntinugwa C, Obah I F, Ike S C, Eme G E, Opara E C, Okolo J C, Nwanyanwu, C E. 2007. In vitro effects of metals and pesticides on dehydrogenase activity in microbial community of cowpea (*Vigna unguiculata*) rhizosphere. *African Journal of Biotechnology* 6: 290-295.
- Omar S A, Abdel-Sater M. A. 2001. Microbial populations and enzyme activities in soil treated with pesticides. *Water Air Soil Pollution* 127: 49-63.
- Panda S, Sahu S K. 2004. Recovery of acetylcholine esterase activity of Drawidawillsi (Oligochaeta) following application of three pesticides to soil. *Chemosphere* 55: 283-290.
- Qian H, Hu B, Wang Z, Xu X, Hong T. 2007. Effects of validamycin on some enzymatic activities in soil. *Environment Monitoring Assessment* 125: 1-8.
- Ramudu A C, Jaffer Mohiddin G, Srinivasulu M, Madakka M, Rangaswamy V. 2011. Impact of fungicides chlorothalonil and propiconazole on microbial activities in groundnut (*Arachis hypogaea* L.) Soils. *ISRN Microbiology* doi:10.5402/2011/623 404.
- Rico A, Sabater C, Castillo M A. 2016. Lethal and sub-lethal effects of five pesticides used in rice farming on the earthworm *Eisenia foetida*. *Ecotoxicology and Environmental Safety* 127: 222-229.
- Romero E, Fernandez-Bayo J, Diaz J M C, Nogales R. 2010. Enzyme activities and diuron persistence in soil amended with vermicompost derived from spent grape marc and treated with urea. *Applied Soil Ecology* 44: 198-204.
- Seghers D, Verthe K, Reheul D, Bulcke R, Siciliano S D, Verstraete W, Top E M. 2003. Effect of long-term herbicide applications on the bacterial community structure and function in an agricultural soil. *FEMS Microbiology Ecology* 46: 139-146.
- Singh G, Wright D. 1999. Effect of herbicides on nodulation, symbiotic nitrogen fixation, growth and yield of pea (*Pisum sativum*). *Journal of Agricultural Science* 133: 21-30.
- Sreenivasa M N, Bagyaraj D J. 1989. Use of pesticides for mass production of vesicular–arbuscular mycorrhizal inoculum. *Plant and Soil* 119: 127-132.
- Stepic S, Hackenberger B K, Velki M, Loncari Z, Hackenberger D K. 2013. Effects of individual and binary-combined commercial insecticides endosulfan, temephos, malathion and pirimiphos-methyl on biomarker responses in earthworm *Eisenia andrei*. *Environmental Toxicology and Pharmacology* 36: 715-723.
- Strandberg M, Scott-Fordsmann J J. 2004. Effects of pendimethalin at lower trophic levels- a review. *Ecotoxicology and Environmental Safety*. 57: 190-201.
- Subash S P, Chand P, Pavithra S, Balaji S J, Pal S. 2017. Pesticide use in Indian agriculture: Trends, market structure and policy issues. Policy brief Technical Report, ICAR-NIAEPR, New Delhi.

- Sukul P. 2006. Enzymes activities and microbial biomass in soil as influenced by metalaxyl residues. *Soil Biology and Biochemistry* 38: 320-326.
- Sumalan R M, Alexa E, Negrea M, Sumalan R L, Doncean A, Pop G. 2010. Effect of glyphosate on the microbial activity of two Romanian soils. *Communication of Agriculture and Applied Biology Sciences* 75: 167-172.
- Sushil S N. 2016. Emerging issues of plant protection in India. *Natural Resource Management: Ecological perspectives*. International Conference, SKUAST, Jammu
- Svensson J, Leonardsson L. 1992. Effects on agricultural pesticides on denitrification in lake sediments. *Proceedings. International Symposium on Environmental Aspects of Pesticide Microbiology*. 55-60 pp.
- Thomas MB. 1999. Ecological approaches and the development of "truly integrated" pest management. *Proceedings of the National Academy of Sciences of the United States of America*. 96: 5944-5951.
- Tu C M. 1995. Effect of 5 insecticides on microbial and enzymatic-activities in sandy soil. *Journal of Environment Science Health* 30: 289-306.
- Vejsadova H, Hrselova H, Prikryl Z, Vancura V. 1989. Effect of different phosphorus and nitrogen levels on development of VA mycorrhiza, rhizobia activity and plant growth. *Agriculture Ecosystem Environment* 29: 429-434.
- Wang C, Zhang Q, Wang F, Liang W. 2016. Toxicological effects of dimethomorph on soil enzymatic activity and soil earthworm (*Eiseniafetida*). *Chemosphere* 169: 316-323.
- Wang MC, Liu YH, Wang Q, Gong M, Hua XM, Pang Y J, Hu S, Yang YH. 2008. Impacts of methamidophos on the biochemical, catabolic, and genetic characteristics of soil microbial communities. *Soil Biology and Biochemistry* 40:778-788.
- Wauchope R D, Yeh S, Linders J B H, Kloskowski R, Tanaka K, Katayama A, Ko`rdel W, Gerstl Z, Lane M, Unsworth J. 2001. Pesticide soil sorption parameters: theory, measurement, uses, limitations and reliability. Poster presented at the IUPAC Congress/ General Assembly.
- Weston LA, Duke S O. 2003. Weed and crop allelopathy. *Critical Reviews in Plant Sciences* 22(3&4): 367-389.
- Wooff D A, Seheult A H, Coolen F P A, Worrall F. 1999. Bayesian discrimination with uncertain covariates for pesticide contamination. In: Barnett, V., Stein, A., Turkman, K. (Eds.), *Statistics for the Environment 4: Pollution Assessment and Control*. John Wiley and Sons, Chichester. 337-353 pp.
- Yao X, Min H, Lu Z, Yuan H. 2006. Influence of acetamiprid on soil enzymatic activities and respiration. *European Journal of Soil Biology* 42: 120-126.
- Zawoznik M S Tomaro M L. 2005. Effect of chlorimuron-ethyl on *Bradyrhizobiumjaponicum* and its symbiosis with soybean. *Pest Management Sci.* 61: 1003-1008.
- Zhang Q, Zhang B, Wang C. 2014. Ecotoxicological effects on the earthworm *Eiseniafetida* following exposure to soil contaminated with imidacloprid. *Environ 4SciPollut Res*. Doi 10.1007/s11356-014-3178-z.
- Zhang W, Ricketts T H, Kremen C, Carney K, Swinton S M. 2007. Ecosystem services and dis-services to agriculture. *Ecological Economics* 64(2): 253-260.

(Manuscript Received: September, 2020; Revised: January, 2021;
Accepted: January, 2021; Online Published: August, 2021)
Online published (Preview) in www.entosocindia.org Ref. No. e20360