



## REPELLENCY OF PLANT ESSENTIAL OILS TO KEY COLEOPTERAN STORED GRAIN INSECTS OF RICE

NAVEENKUMAR B PATIL\*, BASANA GOWDA G, TOTAN ADAK, GURU PIRASANNA PANDI G, MAHENDIRAN ANNAMALAI, P C RATH AND MAYABINI JENA

Crop Protection Division, ICAR-National Rice Research Institute, Cuttack 753006, Odisha, India

\*Email: patil2850@gmail.com (corresponding author)

### ABSTRACT

Laboratory studies were conducted to assess the repellent effects of three essential oils from plants viz., orange, eucalyptus and cinnamon oils against four major coleopteran stored grain insect pests of rice viz., *Sitophilus oryzae*, *Oryzaephilus surinamensis*, *Rhyzopertha dominica*, and *Tribolium castaneum*. The % repellency (PR) and index of repellency (RI) were observed to range from 10 to 100% and 0.00 to 0.90, respectively. Eucalyptus oil @ 5% showed maximum repellent action against *Tribolium castaneum*, registering PR and RI values of 93.33 (F=0.921), 100 (F=1.66), 100 (F=3.772) and 0.07, 0.00 and 0.00, respectively at 3, 6 and 12 hrs after treatment and were found significantly superior over rest of the treatments. Chemical profiling of tested oils through GCMS showed presence of 2-3 chemical constituents amounting to >90 % of total composition of oil. The results highlight the repellency effects of the essential oils and indicate that these can be ecofriendly ones for the post-harvest protection of rice.

**Key words:** Orange, eucalyptus, cinnamon oils, *Sitophilus oryzae*, *Oryzaephilus surinamensis*, *Rhyzopertha dominica*, *Tribolium castaneum*, rice, index of repellency, % repellency, GCMS

In India, stored product insect pests in cereals, pulses and oilseeds cause severe post-harvest losses in the range of 3.9-6.0%, 4.3-6.1% and 2.8-10.1%, respectively (Dhingra, 2016). Amongst these pests, the most important and common are the coleopterans attacking stored rice and their products, viz., *Sitophilus oryzae* L., *Oryzaephilus surinamensis* (L.), *Rhyzopertha dominica* (F.) and *Tribolium castaneum* (Herbst). Management of these depends mostly on the use of fumigants and persistent insecticides. Although effective fumigants like phosphine or methyl bromide (only quarantine treatment) are available to manage stored grain insect pests, but substantial increase in awareness of their ill effects viz. toxicity to non-targets, pesticide residues and environmental pollution is noticed in recent days (Benhalima et al., 2004; Collins et al., 2005 and Haririmoghadam et al., 2011). Aromatic essential oils of plant origin are promising alternative to insecticides in protecting the post-harvest produce and are traditionally been used to kill or repel stored-grain insects (Isman, 2006). These are better alternative to conventional insecticides due to their low mammalian toxicity and high volatility (Shaaya et al., 1997; Li and Zou, 2001). Basically, these are volatile in nature; their secondary metabolites are characterized by a strong aroma and having density lower than water (Bakkali et al., 2008). It has been well established that products of

biological origin are reported to have useful insecticidal compounds against insect pests (Arthur, 1996). Recently, these essential oils have been recognised as pesticides (Isman et al., 2011). Essential oils are considered insecticides because they are selectively bioactive, have little or no harmful effects on non-target organisms and environment (Dong et al., 2004; Kestenholz et al., 2007 and Regnault et al., 2012). In this study, repellent activity of three plant essential oils viz., orange (*Citrus sinensis* L.), eucalyptus (*Eucalyptus obliqua* L'Her) and cinnamon (*Cinnamomum verum*) oils are evaluated against stored grain insect pests of rice viz., *S. oryzae*, *T. castaneum*, *O. surinamensis* and *R. dominica* under laboratory conditions.

### MATERIALS AND METHODS

The samples of *S. oryzae*, *T. castaneum*, *R. dominica* and *O. surinamensis* were collected from rice storage godown of ICAR- National Rice Research Institute (NRI), Cuttack and were maintained in the Grain Entomology laboratory (28± 2°C; 65± 5% RH). Initially, 50 pairs of freshly emerged adults were placed in a jar containing rice grains (0.5 kg). The open end of jars were covered with muslin cloth and allowed for 7 days for mating and oviposition. Then parental stocks were removed and the left over content of each jar (freshly laid eggs and rice grains) were allowed for further

multiplication and insects were collected and reused for subculturing. The subsequent progenies (adults) were used for bioassays. For the bioassay, commercially available eucalyptus, orange and cinnamon oils were obtained from commercial suppliers (NICE Chemicals Private Limited, India). For chemical characterization, the tested plant essential oils (2  $\mu$ l) were dissolved in Hexane (HPLC grade) and were analyzed using GC-MS (Jeol GC mate) system following Thanigaivel et al. (2017). The molecular weight/ formula and structure of the compounds of test materials were ascertained by interpretation on mass spectrum of GC-MS using the database of the National Institute of Standards and Technology (NIST).

‘Area preference method’ in a completely randomized design was used to assess the repellent effects of essential oils (Obeng- Ofori et al., 1998) during 2017-2018. Preparation of test solutions (@ 1, 2 and 5 %) was done by dissolving essential oils in acetone (AR grade). Half cut filter paper (diameter 9.0 cm, Whatman No.1) was dipped in 1.0 ml of the respective test oil while, remaining half was dipped in 1.0 ml acetone which served as control. Both halves were allowed for solvent evaporation. All treatment and control halves were attached together on 9 cm glass petridishes using adhesive tape from bottom of filter paper. Three replicates were maintained for each concentration of oil. Test insects (20 Nos) were released at the centre of each filter paper disc and were then covered and sealed using para films. Petridishes were kept under dark at 26°C and 65 $\pm$  5%RH. The number of adults on the treated and untreated sides were counted at 3, 6, 12 and 24 hr after treatment, with the experiment repeated twice. Effectiveness of plant essential oil was evaluated by Percent Repellency (PR) and Repellency Index (RI) using the formula  $PR (\%) = [(N_c - N_t) / (N_c + N_t)] \times 100$  (where,  $N_c$ - no. of insects in untreated side,  $N_t$ - no. of insects in treated side as per Nerio et al., 2009). Ranges of PR values and their categories used are: 0-0.1%: Class 0; 0.1-20%: Class I; 20.1-40%: Class II; 40.1-60%: Class III; 60.1-80%: Class IV; 80.1-100%: Class V (Tapondjou et al., 2005). Repellency index was calculated by using  $RI = 2G / (G+P)$  as per Mazzonetto (2002), wherein  $G$ = number of insects in treated side and  $P$ = number of insects in untreated side. RI values ranges from 0 to 1 and inversely related with PR values.

## RESULTS AND DISCUSSION

The constituents detected in the study from eucalyptus, cinnamon and orange oils are shown in

Table 1. These reveal that the major compounds of eucalyptus oil were eucalyptol (64.80%),  $\alpha$ -pinene (11.17%),  $\beta$ -pinene (8.19%),  $\gamma$ -terpinene (5.91%),  $\alpha$ -phellandrene (3.88%), terpinen-4-ol (0.72%),  $\alpha$ -terpineol (1.01) and 4-carene (0.51%); beside these, carvacrol,  $\alpha$ -terpinene, 1-epi- $\alpha$ -gurjunene, aromandendrene, alloaromadendrene,  $\alpha$ -farnesene,  $\gamma$ -gurjunene,  $\gamma$ -eudesmol and  $\beta$ -eudesmol were observed. Cinnamon oil resulted in eugenol (82.68%) as the major constituent; in addition, caryophyllene (4.60%), safrole (2.19%), trans-isoeugenol (1.82%), caryophyllene oxide (0.88%), humulene (0.56%), linalool (0.51%) were the other major constituents. Orange oil had D-limonene (83.35%) as the major constituent and cis-limonene oxide (3.31%), trans-2-carene-4-ol (2.08%), tricyclo [4.1.0.0(2,7)] heptanes (0.84%), chrysanthenone (0.73%) and  $\beta$ -myrcene (0.59%) were the other constituents.

The present study reveals that the repellent activity of the essential oils depends on the insect pest and time after treatment, with concentration dependent repellent activity noticed with all the four pests evaluated. Toxicity of eucalyptus oil to coleopteran stored product pests had been attributed to metabolic compounds such as terpenoids and phenolic compounds (Lee et al., 2004; Tapondjou et al., 2005), and its toxicity to lepidopteran agricultural pests is known (Isman, 2006). Numerous plant species had been reported to have repellency properties, contact and fumigant toxicity (Golob et al., 1999). Important constituents of *Cymbopogon* spp., *Ocimum* spp. and *Eucalyptus* spp., and repellent activity of essential oils on insects had been reviewed by Nerio et al. (2010). Essential oils and many other plant extracts are known to possess repellent, insecticidal and ovicidal activities against various stored product insects (Ahmed et al. 1980; Hill and Schoonhoven, 1981; Jilani and Saxena, 1990; Desmarchelier, 1994; Papachristos and Stamopoulos, 2002) and was due to presence of monoterpenoids (Tong and Coats, 2010, Waliwitiya et al., 2005). Due to the high volatile nature of oils these are known to possess repellent and fumigant activities that have pest management significance (Koul, 2004; Konstantopoulou et al., 1992). The present results of chemical profiling of essential oils through GC MS showed presence of 2- 3 chemical constituents amounting to >90 % of total composition of oil which is responsible for repellent activity.

Bioassay against stored grain insect pests revealed increased repellency with concentration of evaluated oils. Repellency Index and percent repellency values

Table 1. Chemical constituents of oils used and their relative composition

Cinnamon oil			Eucalyptus oil			Orange oil		
Compound	Retention time	% area	Compound	Retention time	% area	Compound	Retention time	% area
Benzaldehyde	3.29	0.03	$\alpha$ -Pinene	3.08	11.17	$\alpha$ -Pinene	3.05	0.33
o-Cymene	3.95	0.15	$\beta$ -Pinene	3.51	8.19	$\beta$ -Myrcene	3.54	0.59
Linalool	4.81	0.51	$\alpha$ -Phellandrene	3.76	3.88	D-Limonene	4.08	83.35
4-Carene	6.03	0.24	Eucalyptol	4.11	64.80	$\alpha$ -Ocimene	4.83	0.27
cis-p-Mentha-2,8-dien-1-ol	6.20	0.01	$\gamma$ -Terpinene	4.43	5.91	cis-Limonene oxide	5.33	3.31
3-Phenylpropanol	6.53	0.06	4-Carene	4.73	0.51	1,1,2-trimethyl-Cyclopropane	5.86	0.29
Chavicol	6.79	0.10	Terpinen-4-ol	5.89	0.72	Tricyclo[4.1.0.0(2,7)]heptane	6.17	0.84
Safrole	7.33	2.19	$\alpha$ -Terpineol	6.06	1.01	trans-2-Carene-4-ol	6.42	2.08
Cinnamyl alcohol	7.81	0.06	Carvacrol	7.41	0.06	(-)-Carvone	7.12	0.20
Eugenol	8.48	82.68	2,4-dimethyl-1,3-Cyclopentanedione	7.68	0.21	5-ethylidene-1-methyl-Cycloheptene	7.54	0.66
Caryophyllene	9.18	4.60	$\alpha$ -Terpinene	8.09	0.30	2,5,5-trimethyl-1,3,6-Heptatriene	7.72	0.50
Humulene	9.57	0.56	1-Epi-alpha-gurjunene	8.94	0.50	Chrysanthenone	8.25	0.73
$\beta$ -Patchoulene	10.03	0.11	Aromandendrene	9.33	0.43	8-oxo-cis-Ocimene	8.45	0.35
trans-Isoeugenol	10.33	1.82	Alloaromadendrene	9.59	0.17	Caryophyllene oxide	11.09	0.06
Caryophyllene oxide	11.13	0.88	$\alpha$ -Farnesene	10.71	0.08	Santolina triene	12.27	0.08
$\alpha$ -Farnesene	11.91	0.24	$\gamma$ -Gurjunene	11.09	0.23	Eicosane	17.99	0.06
Coniferol	12.73	0.15	$\gamma$ -eudesmol	11.61	0.22			
Benzyl Benzoate	13.11	3.78	$\beta$ -eudesmol	11.85	0.48			
Total	-	98.17	Total	-	98.87	Total	-	93.7

Concentration (%): % of concentrations based on peak area integration.

ranged from 0.00 to 0.90 and 10 to 100 (Table 2). Against *S. oryzae*, eucalyptus oil @ 5% concentration exhibited significantly higher PR 83.33% (F=4.282) and RI value of 0.17 when tested at 3 hr after treatment; cinnamon oil @ 5% concentration found on par with eucalyptus oil @ 5% with same PR and RI values; at 6, 12 and 24 hrs after treatment, cinnamon oil @ 5% exhibited significantly higher repellency with PR values of 96.67 (F=10.68), 90.00 (F=3.857), and 86.67 (F=3.25) and RI of 0.03, 0.10 and 0.13, respectively. A repellent action of eucalyptus oil against stored grain pests is known earlier. Extracts of eucalyptus leaf prepared in different solvents are known to possess repellent action against *S. oryzae* adults (Lee et al., 2004); similarly repellent potential of leaves of six vegetables species against adults of *S. zeamais* was observed by Procópio et al. (2003) and they reported only leaves of *E. citriodora* (PI = -0.81) and *Capsicum frutescens* (PI = -0.17) have shown the repellency based on Preference Index (PI).

Similarly, against *T. castaneum* eucalyptus oil @ 5% showed maximum repellent action with PR

and RI values of 93.33 (F=0.921), 100 (F=1.66), 100 (F=3.772) and 0.07, 0.00 and 0.00 at 3, 6 and 12 hr after treatment, and were found significantly superior; at 12 hr after treatment orange oil @ 5% had shown repellent action on par with eucalyptus oil @ 5% with PR value of 98.33 and RI value of 0.03; and after 24 hr after treatment orange oil @ 5% exhibited significantly superior repellency (F=20.26) with PR and RI values 100 and 0.00, respectively. Against *R. dominica*, at all time intervals, eucalyptus oil @ 5% has shown maximum repellency; at 3 and 6 hr after treatment, the said oil exhibited repellent action with PR value of 86.67 (F=3.87), 83.33 (F=2.60) and RI value of 0.13, 0.17 and found significantly superior; at 12 hr after treatment, eucalyptus oil @ 2% having PR value of 80.00 (F=10.99) and RI value of 0.20 was found on par with eucalyptus oil @ 5% having PR value of 83.33 RI value of 0.17; same eucalyptus oil @ 2% having PR value of 76.67 (F=12.56) and RI value of 0.23 was found on par with eucalyptus oil @ 5% with PR value of 83.33 (F=12.56) and RI value of 0.17 and orange oil @ 5% having PR value of 83.33 (F=12.56) and RI value of 0.17. All the evaluated oils at different

Table 2. Repellent activity of plant essential oils against major stored grain pests of rice

S. No.	Treatments	3 HAT			6 HAT			12 HAT			24 HAT				
		Adult individuals (%)	PR (%)	RI	Adult individuals (%)	PR (%)	RI	Adult individuals (%)	PR (%)	RI	Adult individuals (%)	PR (%)	RI		
		UT	T	UT	UT	T	UT	UT	T	UT	UT	T			
1	Eucalyptus oil @ 1%	78.33	21.67	56.67 <sup>de</sup>	0.43	76.67	88.33	78.33	21.67	56.67 <sup>e</sup>	0.43	71.67	28.33	43.33 <sup>d</sup>	0.57
2	Eucalyptus oil @ 2%	85.00	15.00	70.00 <sup>bc</sup>	0.30	83.33	16.67	81.67	18.33	66.67 <sup>d</sup>	0.33	78.33	21.67	56.67 <sup>e</sup>	0.43
3	Eucalyptus oil @ 5%	91.67	8.33	83.33 <sup>a</sup>	0.17	91.67	8.33	90.00	10.00	80.00 <sup>b</sup>	0.20	90.00	10.00	80.00 <sup>ab</sup>	0.20
4	Cinnamon oil @ 1%	80.00	20.00	60.00 <sup>de</sup>	0.40	86.67	13.33	88.33	11.67	76.67 <sup>b</sup>	0.23	81.67	18.33	63.33 <sup>c</sup>	0.37
5	Cinnamon oil @ 2%	88.33	11.67	76.67 <sup>ab</sup>	0.23	93.33	6.67	91.67	8.33	83.33 <sup>ab</sup>	0.17	88.33	11.67	76.67 <sup>b</sup>	0.23
6	Cinnamon oil @ 5%	91.67	8.33	83.33 <sup>a</sup>	0.17	98.33	1.67	95.00	5.00	90.00 <sup>a</sup>	0.10	93.33	6.67	86.67 <sup>a</sup>	0.13
7	Orange oil @ 1%	65.00	35.00	30.00 <sup>f</sup>	0.70	66.67	33.33	65.00	35.00	30.00 <sup>e</sup>	0.70	73.33	26.67	46.67 <sup>a</sup>	0.53
8	Orange oil @ 2%	76.67	23.33	53.33 <sup>e</sup>	0.47	73.33	26.67	71.67	28.33	43.33 <sup>d</sup>	0.57	80.00	20.00	60.00 <sup>c</sup>	0.40
9	Orange oil @ 5%	81.67	18.33	63.33 <sup>cd</sup>	0.37	76.67	23.33	88.33	11.67	76.67 <sup>b</sup>	0.23	91.67	8.33	83.33 <sup>ab</sup>	0.17
SEm±		4.08		8.16		3.19		5.09		10.18		4.44		8.89	
CD=0.01		16.62		33.24		12.99		20.73		41.45		18.09		36.18	
1	Eucalyptus oil @ 1%	86.67	13.33	73.33 <sup>bc</sup>	0.27	88.33	11.67	88.33	11.67	76.67 <sup>e</sup>	0.23	86.67	13.33	73.33 <sup>d</sup>	0.27
2	Eucalyptus oil @ 2%	90.00	10.00	80.00 <sup>b</sup>	0.20	93.33	6.67	91.67	8.33	83.33 <sup>b</sup>	0.17	93.33	6.67	86.67 <sup>c</sup>	0.13
3	Eucalyptus oil @ 5%	96.67	3.33	93.33 <sup>a</sup>	0.07	100.00	0.00	100.00	0.00	100.00 <sup>a</sup>	0.00	98.33	1.67	96.67 <sup>ab</sup>	0.03
4	Cinnamon oil @ 1%	76.67	23.33	53.33 <sup>f</sup>	0.47	86.67	13.33	88.33	11.67	76.67 <sup>e</sup>	0.23	96.67	3.33	93.33 <sup>b</sup>	0.07
5	Cinnamon oil @ 2%	81.67	18.33	63.33 <sup>de</sup>	0.37	90.00	10.00	93.33	6.67	86.67 <sup>b</sup>	0.13	98.33	1.67	96.67 <sup>ab</sup>	0.03
6	Cinnamon oil @ 5%	86.67	13.33	73.33 <sup>bc</sup>	0.27	91.67	8.33	98.33	1.67	96.67 <sup>a</sup>	0.00	100.00	0.00	100.00 <sup>a</sup>	0.00
7	Orange oil @ 1%	78.33	21.67	56.67 <sup>ef</sup>	0.43	81.67	18.33	80.00	20.00	60.00 <sup>d</sup>	0.40	91.67	8.33	83.33 <sup>c</sup>	0.17
8	Orange oil @ 2%	81.67	18.33	63.33 <sup>de</sup>	0.37	93.33	6.67	91.67	8.33	83.33 <sup>b</sup>	0.17	96.67	3.33	93.33 <sup>b</sup>	0.07
9	Orange oil @ 5%	85.00	15.00	70.00 <sup>cd</sup>	0.30	96.67	3.33	98.33	1.67	96.67 <sup>a</sup>	0.03	100.00	0.00	100.00 <sup>a</sup>	0.00
SEm±		6.41		12.81		4.23		3.24		6.48		2.94		5.88	
CD=0.01		26.08		52.16		17.22		13.19		26.37		11.97		23.93	
1	Eucalyptus oil @ 1%	81.67	18.33	63.33 <sup>c</sup>	0.37	86.67	13.33	81.67	18.33	63.33 <sup>b</sup>	0.37	76.67	23.33	53.33 <sup>c</sup>	0.47
2	Eucalyptus oil @ 2%	88.33	11.67	76.67 <sup>b</sup>	0.23	88.33	11.67	90.00	10.00	80.00 <sup>a</sup>	0.20	88.33	11.67	76.67 <sup>a</sup>	0.23
3	Eucalyptus oil @ 5%	93.33	6.67	86.67 <sup>a</sup>	0.13	91.67	8.33	91.67	8.33	83.33 <sup>a</sup>	0.17	91.67	8.33	83.33 <sup>a</sup>	0.17
4	Cinnamon oil @ 1%	73.33	26.67	46.67 <sup>de</sup>	0.53	78.33	21.67	80.00	40.00	20.00 <sup>f</sup>	0.80	58.33	41.67	16.67 <sup>e</sup>	0.83
5	Cinnamon oil @ 2%	75.00	25.00	50.00 <sup>de</sup>	0.50	80.00	20.00	83.33	36.67	26.67 <sup>e</sup>	0.73	60.00	40.00	20.00 <sup>e</sup>	0.80
6	Cinnamon oil @ 5%	76.67	23.33	53.33 <sup>d</sup>	0.47	86.67	13.33	86.67	33.33	33.33 <sup>d</sup>	0.67	68.33	31.67	36.67 <sup>d</sup>	0.63
7	Orange oil @ 1%	71.67	28.33	43.33 <sup>e</sup>	0.57	68.33	31.67	75.00	25.00	50.00 <sup>e</sup>	0.50	76.67	23.33	53.33 <sup>c</sup>	0.47
8	Orange oil @ 2%	80.00	20.00	60.00 <sup>c</sup>	0.40	73.33	26.67	81.67	18.33	63.33 <sup>b</sup>	0.37	85.00	15.00	70.00 <sup>b</sup>	0.30
9	Orange oil @ 5%	86.67	13.33	73.33 <sup>b</sup>	0.27	88.33	11.67	88.33	11.67	76.67 <sup>a</sup>	0.23	91.67	8.33	83.33 <sup>a</sup>	0.17
SEm±		3.77		7.54		4.87		3.60		7.20		3.64		7.29	
CD=0.01		15.34		30.67		19.84		14.66		29.31		14.83		29.66	
1	Eucalyptus oil @ 1%	66.67	33.33	33.33 <sup>d</sup>	0.67	63.33	36.67	61.67	38.33	23.33 <sup>cd</sup>	0.77	56.67	43.33	13.33 <sup>d</sup>	0.87
2	Eucalyptus oil @ 2%	71.67	28.33	43.33 <sup>c</sup>	0.57	66.67	33.33	68.33	31.67	36.67 <sup>ab</sup>	0.63	65.00	35.00	30.00 <sup>bc</sup>	0.70
3	Eucalyptus oil @ 5%	80.00	20.00	60.00 <sup>b</sup>	0.40	68.33	31.67	71.67	28.33	43.33 <sup>a</sup>	0.57	68.33	31.67	36.67 <sup>ab</sup>	0.63
4	Cinnamon oil @ 1%	63.33	36.67	26.67 <sup>d</sup>	0.73	61.67	38.33	58.33	41.67	16.67 <sup>de</sup>	0.83	61.67	38.33	23.33 <sup>c</sup>	0.77
5	Cinnamon oil @ 2%	65.00	35.00	30.00 <sup>d</sup>	0.70	70.00	30.00	65.00	35.00	30.00 <sup>de</sup>	0.70	65.00	35.00	30.00 <sup>de</sup>	0.70
6	Cinnamon oil @ 5%	71.67	28.33	43.33 <sup>c</sup>	0.57	81.67	18.33	70.00	30.00	40.00 <sup>a</sup>	0.60	70.00	30.00	40.00 <sup>a</sup>	0.60
7	Orange oil @ 1%	78.33	21.67	56.67 <sup>b</sup>	0.43	65.00	35.00	55.00	45.00	10.00 <sup>e</sup>	0.93	56.67	43.33	13.33 <sup>d</sup>	0.87
8	Orange oil @ 2%	81.67	18.33	63.33 <sup>b</sup>	0.37	70.00	30.00	58.33	41.67	16.67 <sup>de</sup>	0.83	61.67	38.33	23.33 <sup>c</sup>	0.77
9	Orange oil @ 5%	86.67	13.33	73.33 <sup>a</sup>	0.27	85.00	15.00	61.67	38.33	23.33 <sup>cd</sup>	0.77	68.33	31.67	36.67 <sup>ab</sup>	0.63
SEm±		5.30		10.60		5.24		5.44		10.89		6.74		13.47	
CD=0.01		21.57		43.15		21.33		22.16		44.32		27.42		54.84	

PR - % Repellency; RI - Repellence index; HAT- hr after treatment; UT- Untreated area; T- Treated area



concentrations against *O. surinamensis* did not exhibit significant differences.

Thus, the present results reveal the excellent repellency exhibited by eucalyptus oil @ 5% against *T. castaneum* and *R. dominica*. These results are in line with those of earlier ones- viz., Olivero-Verbal et al. (2010) with *T. castaneum* observed that *Eucalyptus citriodora* is a good repellent. Essential oils of *E. kingsmillii* and *E. salmonophloia* tested at four concentrations against female adults of *Tetranychus urticae* Koch indicated that repellency index was neutral at 9 and 17%, whereas, at 23 and 29% it stands as repellent (Haririmoghadam et al., 2011). The essential oil of *E. dundasii* and *E. astringens* against *O. surinamensis* show a more repellent effect corroborating the present results (Khemira et al., 2012). Study of Salvadores et al. (2007) indicated that clove oil was the best repellent against *R. dominica*, *S. oryzae* and *T. castaneum*. Strong repellent and deterrent activity against *T. castaneum* was observed with the leaf extract of *Ocimum viride* (Owsu, 2001); fumigant and repellent effects of oil of *Ocimum gratissimum* and its constituents as a better alternatives to synthetic fumigants against *T. castaneum*, *S. oryzae*, *R. dominica*, *O. surinamensis* and *Callosobruchus chinensis* (Ogendo et al., 2008). Insecticidal and repellent properties of *C. citrinus* against *C. maculatus* were described by Zandi- Sohani et al. (2013) and repellent effects of essential oils of *E. citriodora*, *Lippia origanoides*, *Tagetes lucida* against *S. zeamais* are known earlier (Nerio et al., 2009); also, repellency against *T. castaneum* with *Piper retrofractum* oil had been documented (Tripathi et al. 2000). Wild species of 21 botanical families screened by Pascual-Villalobos (1999) revealed that the family Compositae reveals maximum repellency against *T. castaneum*; likewise, antifeedant and repellent properties of *Cyperus articulatus* against *T. castaneum* are known (Abubakar et al., 2000).

Thus, the current study revealed repellent activity of the essential oils against key stored grain insect pests. Traditionally plant based products have been used to kill or repel stored grain insects since time immemorial, hence the evaluated essential oils could be better alternative to conventional insecticides.

#### ACKNOWLEDGEMENTS

Authors acknowledge the technical and financial support provided by the Director, ICAR-National Rice Research Institute, Cuttack (Odisha); and thank Mr Narendra Biswal and Mr Rabi Bhoi for their assistance.

#### REFERENCES

- Abubakar M S, Abdulrahman E M, Haruna A K. 2000. The repellent and antifeedant properties of *Cyperus articulatus* against *T. castaneum*. *Phytotherapy Research* 14(4): 281-283.
- Ahmed A, Sultana P, Ahmed A. 1980. Comparative efficacy of some indigenous plant materials as repellents against *Sitophilus oryzae* Linn. *Bangladesh Journal of Agricultural Research* (5)2: 31-35.
- Arthur F H. 1996. Grain Protectants: current status and prospects for the future. *Journal of Stored Product Research* 32: 293-302.
- Bakkali F, Averbeck S, Averbeck D, Idaomar M. 2008. Biological effects of essential oils - A review. *Food and Chemical Toxicology* 46: 446-475.
- Benhalima H, Chaudhry M Q, Mills K A, Price N R. 2004. Phosphine resistance in stored-product insects collected from various grain storage facilities in Morocco. *Journal of Stored Product Research* 40: 241-249.
- Collins P J, Daglish G J, Pavic H, Kopitke K A. 2005. Response of mixed-age cultures of phosphine-resistant and susceptible strains of the lesser grain borer, *Rhyzopertha dominica*, to phosphine at arrange of concentrations and exposure periods. *Journal of Stored Product Research* 41(4): 373-385.
- Desmarchelier J M. 1994. Grain protectants: trends and developments. In: Highley E, Wright E J, Banks H J, Champ B R (Eds.), *Stored Product Protection*. Vol. 2. CAB International, Wallingford, UK, pp. 722-728.
- Dhingra D. 2016. Evolution and trends in food grain storage in India. In: Navarro S, Jayas D S, Alagusundaram K, (Eds.) *Proceedings. 10<sup>th</sup> International conference on controlled atmosphere and fumigation in stored products (CAF 2016)*, CAF Permanent Committee Secretariat, Winnipeg, Canada. pp. 47-52.
- Dong Z J, Cheng D Q, Dong S D, Zhou Y H. 2004. Botanical pesticide research and application in tobacco pest control. *Acta Tabacaria Sinica* 10: 42-47.
- Golob P, Moss C, Dales M, Fidgen A, Evans J. 1999. The use of spices and medicinals as bioactive protectants for grains. *FAO Agricultural Services Bulletin* 137.
- Haririmoghadam F, Moharramipour S, Sefidkon F. 2011. Repellent activity and persistence of essential oil from (*Eucalyptus salmonophloia* F. Muell) and (*Eucalyptus kingsmillii* (Mauden) Maiden & Blakely) on two spotted spider mite, (*Tetranychus urticae* Koch). *Iranian Journal of Medicinal and Aromatic Plants* 27(3): 54-65.
- Hill J M, Schoonhoven A V. 1981. The use of vegetable oils in controlling insect infestations in stored grains and pulses. *Recent Advances in Food Science and Technology* 1: 473-481.
- Isman M B. 2006. Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annual Review of Entomology* 51: 45-66.
- Isman M B, Miresmailli S, Machial C. 2011. Commercial opportunities for pesticides based on plant essential oils in agriculture, industry and consumer products. *Phytochemicals Review* 10: 197-204.
- Jilani G, Saxena R C. 1990. Repellent and feeding deterrent effects of turmeric oil, sweet flag oil, neem oil and a neem based insecticide against lesser grain borer. *Journal of Economic Entomology* 83: 629-634.
- Kestenholz C, Stevenson P C, Belmain S R. 2007. Comparative study of field and laboratory evaluations of the ethnobotanical *Cassia sophera* L. (Leguminosae) for bioactivity against the storage pests *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) and *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae). *Journal of Stored Product Research* 43: 79-86.

- Khemira S, Mediouni-Ben Jemâa J, Haouel S, Khouja M L. 2012. Repellent activity of essential oil of *Eucalyptus astringens* against *Rhyzopertha dominica* and *Oryzaephilus surinamensis*. Acta Horticulturae 997: 207-213.
- Konstantopoulou L L, Vassilopoulou L, Mavragani-Tsipidou P, Scouras Z G. 1992. Insecticidal effects of essential oils. A study of the effects of essential oils extracted from eleven Greek aromatic plants on *Drosophila auraria*. Experientia 48: 616-619.
- Koul O. 2004. Biological activity of volatile di-n-propyl disulfide from seeds of neem, *Azadirachta indica* (Meliaceae), to two species of stored grain pests, *Sitophilus oryzae* (L.) and *Tribolium castaneum* (Herbst). Journal of Economic Entomology 97: 1142-1147.
- Lee B H, Annis P C, Tumaalii F, Lee S E. 2004. Fumigant toxicity of *Eucalyptus blakelyi* and *Melaleuca fulgens* essential oils and 1, 8-cineole against different development stages of the rice weevil *Sitophilus oryzae*. Phytoparasitica 32: 498-506.
- Li Y S, Zou H Y. 2001. Insecticidal activity of extracts from *Eupatorium adenophorum* against four stored grain insects. Entomological Knowledge 38: 214-216.
- Mazzonetto F. 2002. Efeito de genótipos de feijoeiro e de pós origem vegetal sobre *Zabrotes subfasciatus* (Boh.) e *Acanthoscelides obtectus* (Say) (Col. Bruchidae). Tesis Doctor en Ciencias. Universidad de Sao Paulo, Piracicaba, Sao Paulo, Brasil. 134 pp.
- Nerio L S, Olivero-Verbal J, Stashenko E E. 2009. Repellent activities of essential oils from seven aromatic plants grown in Colombia against *Sitophilus zeamais* Motschulsky (Coleoptera). Journal of Stored Product Research 45: 212-214.
- Nerio L S, Olivero-Verbel J, Stashenko E. 2010. Repellent activity of essential oils: a review. Bioresource Technology 101: 372-378.
- Obeng-ofori D, Reichmuth C H, Bekele A J, Hassanali A. 1998. Toxicity and protectant potential of camphor, a major component of essential oil of *Ocimum kilimandscharicum*, against four stored product beetles. International Journal of Pest Management 44: 203-209.
- Ogendo J O, Kostyukovski M, Ravid U, Matasyoh J C, Deng A L, Omolo E O, Kariuki S T, Shaaya E. 2008. Bioactivity of *Ocimum gratissimum* L. oil and two of its constituents against five insect pest attacking stored food products. Journal of Stored Product Research 44: 328-334.
- Olivero-Verbel J, Nerio L S, Stashenko E E. 2010. Bioactivity against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) of *Cymbopogon citratus* and *Eucalyptus citriodora* essential oils grown in Colombia. Pest Management Science 66: 664-665.
- Owsu E O. 2001. Effect of some Ghanaian plant components on control of two stored product insect pests of cereals. Journal of stored Product Research 37(1): 85-91.
- Papachristos D P, Stamopoulos D C. 2002. Repellent, toxic and reproduction inhibitory effects of essential oil vapours on *Acanthoscelides obtectus* (Say). Journal of Stored Product Research 38: 117-128.
- Pascual-Villalobos M J. 1999. Anti-insect activity of plant extracts from the wild flora in Southeastern Spain. Biochemical Systematics and Ecology 27 (1): 1-10.
- Procopio S O, Vendramim J D, Ribeiro Júnior J I, Santos J B. 2003. Bioatividade de diversos pós de origem vegetal em relação à *Sitophilus zeamais* Mots. (Coleoptera: Curculionidae). Ciencia e Agrotecnologia 27: 1231-1236.
- Regnault-Roger C, Vincent C, Amasson J T. 2012. Essential oils in insect control: low-risk products in a high-stakes world. Annual Review of Entomology 57: 405-425.
- Salvadores Y U, Silva G A, Tapia M V, Hepp R G. 2007. Spices powders for the control of maize weevil, *Sitophilus zeamais* Motschulsky in stored wheat. Chilean Journal of Agricultural Research 67: 147-154
- Shaaya E, Kostjukovski M, Eilberg J, Sukprakarn C. 1997. Plant oils as fumigants and contact insecticides for the control of stored-product insects. Journal of Stored Product Research 33: 7-15.
- Tapondjou A L, Adler C, Fontem D A, Bouda H, Reichmut C. 2005. Bioactivities of cymol and essential oils of *Cupressus sempervirens* and *Eucalyptus saligna* against *Sitophilus zeamais* (Motschulsky) and *Tribolium confusum* (du Val). Journal of Stored Product Research 41: 91-102.
- Thanigaivel A, Vasantha Srinivasan P, Senthil Nathan S, Edwin E, Ponsankar A, Chellappandian M, Selin Rani S, Lija Escaline J, Kalaivani K. 2017. Impact of *Terminalia chebula* Retz. against *Aedes aegypti* L. and non-target aquatic predatory insects. Ecotoxicology Environmental Safety. 137: 210-217.
- Tong F, Coats J R. 2010. Effects of monoterpene insecticides on (3H)-TBOB binding in house fly GABA receptor and 36 Cl- uptake in American cockroach ventral nerve cord. Pesticide Biochemistry and Physiology 98 (3): 317-324.
- Tripathi A K, Prajapati V, Aggarwal K K, Khanuja S P S, Kumar S. 2000. Repellency and toxicity of oil from *Artemisia annua* to certain stored product beetles. Journal of Economic Entomology 93 (1): 43-47.
- Waliwitiya R, Isman M B, Vernon R S, Riseman A. 2005. Insecticidal activity of selected monoterpenoids and rosemary oil to *Agriotes obscurus*. (Coleoptera: Elateridae). Journal of Economic Entomology 98(5): 1560-1565.
- Zandi Sohani N, Hojjati M, Carbonell Barrachina. 2013. Insecticidal and repellent activities of the essential oil of *Callistemon citrinus* (Myrtaceae) against pulse beetle *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). Neotropical Entomology 42 (1): 89-94.

(Manuscript Received: September, 2020; Revised: January, 2021;  
Accepted: January, 2021; Online Published: August, 2021)  
Online published (Preview) in www.entosocindia.org Ref. No. e20388