



DIVERSITY AND POPULATION DYNAMICS OF SPIDERS IN AGROECOSYSTEMS

S RAGHUL* AND K KUMAR

Department of Agricultural Entomology, Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal 609603, Puducherry, India

*Email: raghulsubasekaran@gmail.com (corresponding author)

ABSTRACT

The biodiversity of spiders in agroecosystem was studied at the Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal during kharif (2019) and rabi (2019-20). The spiders were collected at weekly intervals using in situ counts, net sweeping, pitfall traps and litter sampling. A total of 30 species under 22 genera, 15 families were observed. Biodiversity indices viz., Shannon-Weiner Index (2.809- kharif and 2.766- rabi), Simpson Index (0.926- kharif and 0.909- rabi), Margalef Index (4.135- kharif and 4.22- rabi) and Pielou's Index (0.104- kharif and 0.095- rabi) were computed. Regression with weather parameters during kharif 2019, were non-significant for *Thomisus* sp. (0.107), *Pardosa sumatrana* Thorell (0.146), *Oxyopes javanus* Thorell (0.190), *Tetragnatha javana* Thorell (0.213) and *Tetragnatha mandibulata* Walckenaer (0.347); and during rabi 2019-20, for *T. javana* (0.516), *Argiope anasuja* Thorell (0.619) showed significance and *O. javanus* (0.192), *Lycosa bistrata* Gravely (0.370), *T. mandibulata* (0.437), these values were non-significant.

Key words: Karaikal, kharif, rabi, agroecosystem, spiders, biodiversity indices, population dynamics, correlation, regression, dominance, species richness, abundance

Arachnids are the largest and successful group of chelicerates. Among the arachnids, the order Araneae is the largest group (Thompson, 2015). They are the most diversified group amongst invertebrates with 48,901 species under 4,184 genera and 128 families (WSC, 2020). About 1,909 species belonging to 488 genera and 64 families are from India (WSC, 2019). Spiders play an important predatory role in agroecosystem by lowering insect densities, as well as stabilizing pest populations (Saranya et al., 2019). Spiders have been evidenced as bio-indicators in environmental habitats that could be helpful for conservation purposes (Gerlach et al., 2013). Benamu (2020) stated that studies on spider's diversity in agroecosystem have increased, demonstrating their potential to be used as biological control agents in IPM, and it can reduce the indiscriminate use of pesticides. Spiders are very sensitive to the variations in abiotic conditions, and Pitilin et al. (2019) observed that spiders influence the pest populations in the field and these are also influenced by the weather factors. Hence, it is essential to know the population dynamics in relation to weather factors and the present study evaluates the same under agroecosystem at Karaikal, U T of Puducherry.

MATERIALS AND METHODS

The present study was conducted at the Eastern farm at (10°55'N, 79°49'E, 8 masl) of Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal, U T of Puducherry. Spider fauna were

collected in the early hours (08:00- 10:00 hr.), in the afternoon (14.00- 15.00 hr.) and in late evening (17:00- 18:00 hr.) at weekly intervals using different methods viz., in situ count, net sweeping (Pandit and Pai, 2017), pitfall trap and litter sampling from crops like cotton, maize, rice, ragi and pigeon pea. Pitfall traps (5 no.) were placed at five random spots with 2- 3 drops of liquid soap as trapping fluid and specimens were collected next morning (Bukhari et al., 2012). Litter sampling was done by manual searching of spiders under the leaf litters at weekly intervals (Jose et al., 2018). Following Engelmann (1978), the families were distinguished with the relative abundance as- subprecedent, recedent, subdominant, dominant and eudominant. The collected specimens were killed and preserved in glass vials containing 70% alcohol before labeling. Tikader and Bal (1981) was used for the identification of species.

Diversity indices like Shannon-Wiener index (Hughes, 1978), Simpson's diversity index (Simpson, 1949), Margalef index (Margalef, 1958) and Pielou's evenness index (Pielou, 1966) were computed using standard methodology (https://www.alyoung.com/labs/biodiversity_calculator.html). The weather factors viz., maximum and minimum temperature, morning and evening relative humidity (RH), bright sunshine hours and rainfall obtained were correlated with the occurrence of spiders with correlation coefficients and regression ($p \leq 0.05$ (*) and ≤ 0.01 (**)).

RESULTS AND DISCUSSION

The study revealed a total of 30 spider species with 1,366 individuals under 22 genera and 15 families; Araneidae was the richest with 5 species followed by Oxyopidae with 4 species, Salticidae, Thomisidae and Tetragnathidae (3 species each), Lycosidae and Sparassidae (2 species each), and families Clubionidae, Corinnidae, Eutichuridae, Gnaphosidae, Philodromidae, Pisauridae, Theridiidae and Zodariidae with 1 species each (Table 1). Ambily and Antony (2016) stated that family Araneidae dominated the agroecosystem of Ernakulum, district of Kerala with 8 species. During kharif 2019 and rabi 2019-20 all the families except Corinnidae (*Castianeira zetes*) (kharif and absent during rabi), Theridiidae (*Argyrodes argentatus*) (kharif absent; rabi present), Thomisidae (*Platythomisus sudeepi*) (kharif absent; rabi present), *Tmarus fasciolatus* (kharif absent; rabi present) were observed. The relative abundance of spider fauna during kharif 2019 and rabi 2019-20 revealed that family Tetragnathidae was dominant (21.81%) followed by Oxyopidae (21.53%) and Araneidae (15.23%); Lycosidae (10.03%), Salticidae (7.68%), Thomisidae (5.48%) and Eutichuridae (4.69%) were subdominant. Pisauridae (0.44%), Theridiidae (0.15%) and Corinnidae (0.07%) were

subrecedent; Philodromidae (3.66%), Gnaphosidae (3.00%), Clubionidae (2.71%), Zodariidae (2.20%) and Sparassidae (1.31%) were grouped as recedent (Table 2). Sebastian et al. (2005) stated that the Tetragnathidae was found to have high relative abundance in the irrigated rice ecosystem of Central Kerala. Ranjini (2016) also observed that Tetragnathidae (50%) was the dominant in the rice ecosystem of Palakkad district.

The biodiversity analysis indices revealed that the following viz., Shannon-Weiner Diversity Index (H') (was 2.809 in kharif and 2.766 in rabi), Simpson Dominance Index (D) was (0.926 in kharif and 0.909 in rabi), Margalef Richness Index (α) was (4.135 in kharif and 4.22 in rabi) and Pielou's Evenness Index (E1) was (0.104 in kharif and 0.095 in rabi). These indicate that the species diversity and evenness indices during kharif was more abundant compared to that of rabi; and species richness were more or less equal and exhibited a similar diversification in both the seasons. Anitha and Vijay (2016) reported that the Shannon Wiener Index (H') value was 1.53 and 1.81, Simpson Index (D) value was 0.29 and 0.19, Margelef species richness index value was 1.00 and 1.10, Pielou's Evenness Index (E1) was 0.69 and 0.76 in kharif and rabi, respectively in the rice ecosystem of Rajendranagar, Telangana.

Table 1. List of spiders observed in agroecosystems (Karaikal)*

1.	Araneidae	Orb web spiders	<i>Argiope catenulata</i> (Doleschall) <i>Argiope anasuja</i> (Thorell) <i>Larinia chloris</i> (Audouin) <i>Larinia</i> sp. <i>Neoscona theisi</i> (Walckenaer)
2.	Clubionidae	Sac spiders	<i>Clubiona drassodes</i> (O. Pickard-Cambridge)
3.	Corinnidae	Ant mimic sac spiders	<i>Castianeira zetes</i> (Simon)
4.	Eutichuridae	Yellow sac spiders	<i>Cheiracanthium melanostomum</i> (Thorell)
5.	Gnaphosidae	Ground spiders	<i>Zelotes</i> sp.
6.	Lycosidae	Wolf spiders	<i>Lycosa bistrata</i> (Gravely) <i>Pardosa sumatrana</i> (Thorell)
7.	Oxyopidae	Lynx spiders	<i>Oxyopes javanus</i> (Thorell) <i>Oxyopes shweta</i> (Tikader) <i>Oxyopes sunandae</i> (Tikader) <i>Peucetia viridana</i> (Stoliczka)
8.	Philodromidae	Running crab spiders	<i>Thanatus</i> sp.
9.	Pisauridae	Nursery web spiders	<i>Dolomedes fimbriatus</i> (Clerck)
10.	Salticidae	Jumping spiders	<i>Carrhotus viduus</i> (Koch) <i>Carrhotus sannio</i> (Thorell) <i>Hyllus semicupreus</i> (Simon)
11.	Sparassidae	Giant crab spiders	<i>Olios lamarcki</i> (Latrielle) <i>Olios milleti</i> (Pocock)
12.	Tetragnathidae	Long jawed spiders	<i>Tetragnatha mandibulata</i> (Walckenaer) <i>Tetragnatha javana</i> (Thorell) <i>Tetragnatha viridorufa</i> (Gravely)
13.	Theridiidae	Comb-footed spiders	<i>Argyrodes argentatus</i> (O. Pickard-Cambridge)
14.	Thomisidae	Crab spider	<i>Platythomisus sudeepi</i> (Biswas) <i>Thomisus</i> sp. <i>Tmarus fasciolatus</i> (Simon)
15.	Zodariidae	Ant spider	<i>Malinella</i> sp.

Table 2. Relative abundance of spider fauna in agroecosystem (July 2019 to February 2020)

S. No.	Family	No. of Genera	No. of species	Nos.	Relative abundance (%)		Dominance
					kharif	rabi	
1	Araneidae	3	5	208	9.5	5.73	Dominant
2	Clubionidae	1	1	37	1.45	1.26	Recedent
3	Corinnidae	1	1	1	0.07	0	Subrecedent
4	Eutichuridae	1	1	64	2.65	2.04	Subdominant
5	Gnaphosidae	1	1	41	1.6	1.4	Recedent
6	Lycosidae	2	2	137	6.28	3.75	Subdominant
7	Oxyopidae	2	4	294	12.03	9.5	Dominant
8	Philodromidae	1	1	50	2.1	1.56	Recedent
9	Pisauridae	1	1	6	0.23	0.21	Subrecedent
10	Salticidae	2	3	105	4.29	3.39	Subdominant
11	Sparassidae	1	2	18	0.92	0.39	Recedent
12	Tetragnathidae	1	3	298	15.2	6.61	Dominant
13	Theridiidae	1	1	2	0	0.15	Subrecedent
14	Thomisidae	3	3	75	3.38	2.1	Subdominant
15	Zodariidae	1	1	30	1.6	0.6	Recedent
Total		22	30	1366			

* RA below (1.3)- subrecedent; (1.3 - 3.9) - Recedent; (4 - 12.4) - Subdominant; (12.5 - 39.9) - Dominant and (40 - 100) - Eudominant (Engelmann, 1978).

Correlation with weather parameters during kharif 2019 revealed that *O. javanus* showed negatively significant correlation with bright sunshine (-0.06); similarly *P. sumatrana* with evening RH, and bright sunshine (-0.02, -0.08), and positively significant (0.04) with total rainfall; *T. javana* showed a positively significant one with maximum temperature and total rainfall (0.01, 0.02) and a negatively significant one with morning and evening RH (-0.01, -0.03); *Thomisus* sp. showed a positively significant one with minimum temperature, bright sunshine (0.05, 0.02) and a negative one with morning and evening RH, and total rainfall (-0.07, -0.09, -0.03). There existed a non-significant regression with the weather parameters for *O. javanus* (0.190), *T. mandibulata* (0.347), *T. javana* (0.213), *P. sumatrana* (0.146) and *Thomisus* sp. (0.107). Correlation with weather parameters during rabi 2019-20, showed *L. bistrriata* being positively and significantly correlated with total rainfall (0.01); *O. javanus* with a positively significant (0.01) one with minimum temperature and a negatively significant one with total rainfall (-0.07); with *T. javana*, a positively significant (0.08, 0.01, 0.08) correlation with morning and evening RH and total rainfall; *T. mandibulata* showed a positively significant correlation with minimum temperature, and morning and evening RH (0.04, 0.02 and 0.50).

Yadav et al. (2017) reported that maximum RH had a significant and positive impact on the population of the *Oxyopes* sp. in rice agroecosystem of Bihar; Sidar et al. (2017) reported a non-significant positive correlation with maximum (0.074) and minimum temperature (0.28), morning (0.27) and evening RH

(0.15), whereas rainfall (-0.20), wind velocity (-0.39) and sun shine hours (-0.14) showed a non-significant negative correlation with maize in Chhattisgarh. Patel et al. (2020) reported that morning and evening RH and rainfall exhibited a positive correlation with spiders in cotton. In Telangana, the abundance of spiders revealed a positive correlation with RH and a negative one with temperature and rainfall in rice (Laxman et al., 2016).

ACKNOWLEDGEMENTS

The authors thank Dr M Sampath Kumar, NBAIR, Bengaluru, Dr A V Sudhikumar, Centre for Animal Taxonomy and Ecology, Christ College, Kerala and Dr D A Prajapati, Gujarat Ecological Education and Research Foundation, Gujarat for their guidance in identification of spiders. I thank PAJANCOA and RI Society, Karaikal, U T of Puducherry for permission to conduct the study.

REFERENCES

- Ambily C B, Antony A. 2016. Diversity and distribution of spiders in agroecosystem of Ernakulum district, Kerala. *Journal of Zoological Studies* 3(5): 73-77.
- Anitha G, Vijay J. 2016. Quantification of the abundance and diversity of predatory spiders in rice ecosystem of Rajendranagar, Telangana, India. *Journal of Applied and Natural Science* 8(2): 1010-1014.
- Benamu M A. 2020. The importance of spider diversity in agroecosystems and the effect of pesticides. *Global Journal of Ecology* 5(1): 60-61.
- Bukhari M, Naeem M M, Rehman K, Andleeb S. 2012. Determination of diversity indices of Araneidae fauna captured from guava orchards by pitfall trapping method at Gujranwala. *Pakistan Journal of Entomology* 5: 133-136.
- Engelmann H D. 1978. Zur Dominanzklassi kazierung von Bodenarthropoden. *Pedobiologia* 18: 378-380.

Table 3. Population dynamics of spider species (kharif 2019, rabi 2019-20)

Spider species	Kharif 2019				Rabi 2019-20			
	Temperature (°C)		Relative humidity (%)		Temperature (°C)		Relative humidity (%)	
	Maximum	Minimum	Morning	Evening	Maximum	Minimum	Morning	Evening
<i>Oxyopes javanus</i>	-0.34	-0.32	0.39	0.32	0.60	0.46	-0.68	-0.38
	Y = -19.92-1.18X ₁ +1.40X ₂ +0.62X ₃ -0.31X ₄ -0.22X ₅ -0.06X ₆		(0.190 NS)		Y = 65.11-0.19X ₁ +0.89X ₂ -0.85X ₃ -0.13X ₄ +1.18X ₅ +0.14X ₆		(0.619*)	
<i>Pardosa sumatrana</i>	0.12	0.13	0.02	-0.02**	-0.44	-0.19	0.26	0.22
	Y = -50.09+0.90X ₁ +0.03X ₂ +0.21X ₃ +0.08X ₄ -0.18X ₅ -0.04X ₆		(0.146 NS)		Y = 22.01-1.77X ₁ +1.05X ₂ +0.13X ₃ +0.02X ₄ -0.29X ₅ -0.16X ₆		(0.370 NS)	
<i>Tetragnatha javana</i>	0.01**	0.12	-0.01**	-0.03**	0.11	0.01**	0.22	-0.14
	Y = 2.80-1.83X ₁ +2.29X ₂ +0.35X ₃ +0.34X ₄ -0.71X ₅ -0.05X ₆		(0.213 NS)		Y = -137.3+0.25X ₁ +1.29X ₂ +1.34X ₃ -0.29X ₄ +0.89X ₅ +0.11X ₆		(0.192 NS)	
<i>Tetragnatha mandibulata</i>	-0.28	-0.24	0.31	0.39	0.26	0.41	0.08*	0.01**
	Y = -33.62+3.64X ₁ -3.38X ₂ -0.53X ₃ -0.86X ₄ -1.02X ₅ -0.07X ₆		(0.347 NS)		Y = -56.18-0.71X ₁ +2.12X ₂ +0.84X ₃ -0.51X ₄ -1.41X ₅ -0.04X ₆		(0.516*)	
<i>Thomisus sp.</i>	0.11	0.05*	-0.07*	-0.09*	-0.16	0.04**	0.02**	0.50*
	Y = -9.31+0.71X ₁ -0.60X ₂ -0.03X ₃ +0.088X ₄ -0.01X ₅ -0.03X ₆		(0.107 NS)		Y = -53.36+3.54X ₁ -3.24X ₂ -0.84X ₃ +1.31X ₄ +0.96X ₅ -0.01X ₆		(0.437 NS)	

* = Significant at p=0.05; ** at p=0.01; NS = Not significant; X₁ = Maximum temperature; X₂ = Minimum temperature; X₃ = Morning relative humidity; X₄ = Evening relative humidity; X₅ = Bright sunshine hours; X₆ = Rainfall

Gerlach J, Samways M, Pryke J. 2013. Terrestrial invertebrates as bioindicators an overview of available taxonomic groups. Journal of Insect Conservation 17: 831-850.

Hughes B D. 1978. The influence of factors other than pollution on the value of Shannon's diversity index for benthic macro-invertebrates in streams. Water Research 12: 359-364.

Jose A C, Sudhin P P, Prasad P M, Sreejith K A. 2018. Spider diversity in Kavvayi river basin, Kerala, Southern India. Current World Environment 13(1): 100-112.

Laxman P, Kiranmai K, Thiruthathi U, Sammaiah C. 2016. Impact of weather factors on predatory spiders in Bt and non Bt-cotton fields of Warangal, Telangana. Biolife 4: 386-391.

Margalef R D. 1958. Information theory in ecology. General Systems 3: 36-71.

Pandit R, Pai I K. 2017. Spiders of Taleigao Plateau, Goa, India. Journal of Environmental Science and Public Health 1(4): 240-252.

Patel Y, Patel P, Sharma R. 2020. Path analysis of abiotic factors affects the population dynamics of spider, *Oxyopes* sp. in cotton. Journal of Entomology and Zoology Studies 8(5): 1722-1726.

Pielou E C. 1966. The measurement of diversity in different types of biological collections. Journal of Threatened Biology 13: 131-144.

Pitilin R B, Prado J, Brescovit A D, Buschini M L T. 2019. Climatic conditions drive the abundance and diversity of spider's community in an atlantic forest fragment Oecologia Australis 23(1): 39-55.

Ranjini S. 2016. Preliminary studies on the spider fauna in selected agro-ecosystems of Kozhinjampara, Palakkad, Kerala. International Journal of Scientific Research 7(10): 13740-13743.

Saranya V S L, Samiayyan K, Prema M S. 2019. Diversity of predatory spider fauna in maize ecosystem. Journal of Biological Control 33(1): 27-35.

Sebastian P A, Sudhikumar A V, Mathew M V, Sunish E. 2005. Diversity of spiders (Araneae) in the Western Ghats. Ashoka Trust Research in Ecology, Bangalore. 235-247 pp.

Sidar Y K, Deole S, Nirmal A, Gajbhiye R K, Bisen M S. 2017. A study on the seasonal distribution of spider fauna in the maize field at Raipur, Chhattisgarh. Journal of Entomology and Zoology Studies 5(2): 1105-1108.

Simpson E H. 1949. Measurement of diversity. Nature 163: 688-696.

Thompson J L. 2015. Spiders, scorpions, centipedes and mites. Elsevier Division. The Commonwealth and International Library, Turkey. 296 pp.

Tikader B K, Bal A. 1981. Studies on orb-weaving spiders of the genera *Neoscona* (Simon) and *Araneus* (Clerck) of the family Araneidae from India. Records of Zoological Survey of India 24: 1-60.

World Spider Catalog. 2019. Version 20.0. Natural History Museum Bern. <http://wsc.nmbe.ch/families>.

World Spider Catalog. 2020. Version 20.0. Natural History Museum Bern. <https://wsc.nmbe.ch/families>.

Yadav M, Goswami T N, Ray S N. 2017. Study of seasonal variations in population dynamics of spider families in paddy agroecosystem, Sabour, Bhagalpur, Bihar, India. International Journal of Agricultural Science and Research 7(4): 479-484.

(Manuscript Received: November, 2020;
 Revised: January, 2021;
 Accepted: January, 2021; Online Published:
 August, 2021)
 Online published (Preview) in www.entosocindia.org
 Ref. No. e20397