



## EFFECT OF FARMING SYSTEMS ON DIVERSITY AND SEASONAL ABUNDANCE OF INSECT PESTS AND THEIR NATURAL ENEMIES IN CAULIFLOWER

ABDUL WAKIL BARAKZAI\*, RAJESHWAR SINGH CHANDEL, PREM LAL SHARMA, SUBHASH CHANDER VERMA, MANEESH PAL SINGH AND PANMA YANKIT

Department of Entomology, Dr Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan 173230, Himachal Pradesh, India

\*Email: abdulwakilbb@gmail.com (corresponding author)

### ABSTRACT

A study was conducted on effect of two farming systems namely zero budget natural farming (ZBNF) and conventional farming (CF) on diversity and seasonal abundance of insect pests of cauliflower and their natural enemies during 2018-19 and 2019-20. Studies revealed that ZBNF system harboured less pest diversity and attracted more natural enemies as compared to the CF system. Natural enemy activity and % parasitisation were maximum in ZBNF system as compared to CF system. The results indicate that indigenous ZBNF formulations and intercropping have a positive effect on the population of natural enemies and repelled the insect pests much better as compared to CF system.

**Key words:** Zero budget natural farming, conventional farming, insect pests, natural enemies, cauliflower, repellency, parasitisation, natural enemy activity

Cauliflower *Brassica oleraceae* var. *botrytis* L., is one of the common vegetables, and introduced in India in 1822 from England (Chatterjee et al., 1986). It is currently grown throughout the country. Cauliflower is attacked by various insect pests (Gaikwad et al., 2018; Raja et al., 2014). A large number of insecticides have been recommended and used to control these. These insecticides have their own harmful effects, including their negative impacts on natural enemies. Agroecosystem is not self-regulated as compared to others due to monocropping, and extra efforts are required to control the pests. Natural enemies play an effective role in IPM (Mandal and Patnaik, 2008). Intercropping, organic amendments generally harbour more natural enemies. Farming systems have great impact on insect pests and natural enemies, and organic practices increase the natural enemies (Gallo and Pekar, 1999). Such comparisons are necessary, as agroecosystems are with increased pest diversity and reduced natural enemies (Yadav 1989; Meena et al., 2002). The effect of organic farming system on natural enemies is required to be brought out and there are many studies. Recently a new system of farming called as zero budget natural farming (ZBNF) has been practiced. However, the studies on the impact of this system on the diversity and abundance of insect pests and their natural enemies are lacking. Natural enemies play key role in insect pest control (Liu et al., 2000). Hence, the current study to study the effects of

ZBNF in comparison to conventional farming (CF) in cauliflower.

### MATERIALS AND METHODS

The study was carried out at the experimental farms of Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh (1300 masl). Cauliflower (PSBK-1) was raised following standard package of practices for ZBNF and CF separately. In ZBNF, the main crop was intercropped with pea (P89), coriander (local selection) and mustard (trap crop). In CF cauliflower was raised as a sole crop, with plot size of 9x 4.5 m. For fertilization and pest management, indigenous plant and cow dung-urine based formulations were used. Jeevamrit, Ghanjeevamrit and Beejamrit were used as fertilizers, while Darekastra @ 500 l ha<sup>-1</sup>, Bramhastra @ 3% and Agniastra @ 3% were used for pest management in ZBNF (Palekar, 2016). In CF system crop was raised following all the standard package of practices recommended by the University. The fertilizers used were urea @ 300 kg, SSP @ 675 kg and MOP @ 85kg/ha. The pests and diseases were controlled by need based applications of Captan (75%WP), malathion (50%EC) and imidacloprid (17.8%SL).

The diversity and seasonal abundance of insect pests were monitored at weekly intervals from nursery stage to final harvest. The data on the number of species

of pests and their natural enemies and the density of each species were observed. The population density of aphids, whitefly, caterpillars and beetles were counted on plant basis from 50 randomly selected plants in each case. The natural enemies viz. coccinellids and syrphid flies' density were observed by directly counting the beetles and maggots/ plant; in case of parasitized aphids, the parasitized ones were collected from 50 randomly selected plants and kept in the laboratory for emergence of adult parasitoids. The parasitized larvae of diamond back moth and cabbage butterfly were collected from sampling plants and kept in the laboratory for the emergence of parasitoids. The diversity of insect pests and natural enemies were analysed through Simpson Index (Simpson, 1949) and Shannon- Wiener Index (Shannon, 1948). For evaluating the seasonal abundance, weekly data collected were statistically analysed using t-Test by OP-STAT software (Sheoran et al., 1998).

## RESULTS AND DISCUSSION

Diversity of insect pests and natural enemies: During 2018-19, six species of insect pests (under 6 genera and families each) were observed on cauliflower crop under ZBNF and CF systems. The pests include: the cabbage aphid *Brevicoryne brassicae* (L.); flea beetle *Phyllotreta* sp; greenhouse whitefly *Trialeurodes vaporariorum* (Westwood); diamond back moth *Plutella xylostella* (L.); cabbage head borer *Hellula undalis* (F.) and cabbage butterfly *Pieris brassicae* (L.). During 2019-20, in ZBNF system the diversity and species composition remained the same as that of previous year. In CF system, in addition to the above insect pests cabbage semilooper *Thysanoplusia orichalcea* (F.) was observed. As regards natural enemies, seven species were found associated with insect pests under ZBNF system. Out of these, five were predators and two parasitoids. Among predators, two coccinellids i.e. *Coccinella septempunctata* (L.) and *Hippodamia*

*variegata* (Goeze); and three syrphids i.e. *Episyrphus balteatus* (De Geer), *Eupeodes frequens* (Matsumura) and *Metasyrphus confrator* (Wiedemann) were observed. The parasitoids observed were *Diaeretiella rapae* (Mc'Intosh) parasitizing the cabbage aphid, and *Diadegma semiclausum* (Hellen) parasitizing the larvae of *P. xylostella*.

During 2019-20 in ZBNF system, eight species of natural enemies were observed, all the species observed in the previous year were found along with the parasitoid *Cotesia glomerata* (L.) parasitizing the larvae of butterfly. In cauliflower grown under the conventional farming system, six species of natural enemies were observed during both the years of study. Five of them were predators and one was parasitoid. Among the five predators, two coccinellids i.e. *C. septempunctata* and *H. variegata*; and three syrphids i.e. *E. balteatus*, *E. frequens* and *M. confrator* were observed. The parasitoid observed was *D. rapae* parasitizing the cabbage aphid. The insect pest diversity indices revealed that CF based cauliflower ecosystem was more diverse than ZBNF system, while, the natural enemies' diversity indices revealed that ZBNF based system was more diverse (Table 1). This is possibly because of higher crop diversity and non-application of insecticides in the ZBNF system.

Seasonal abundance: The cabbage aphid appeared during the last week of October, 2018 (44<sup>th</sup> standard week (SW)- 2.74± 0.30 and 4.18± 0.40 aphids plant<sup>-1</sup> in ZBNF and CF systems, respectively during 2018-19; and reached peak incidence of 23.40± 4.00 and 27.74± 5.09 aphids plant<sup>-1</sup>) during 6<sup>th</sup> and 5<sup>th</sup> SW in ZBNF and CF systems, respectively. In 2019-20, the aphids appeared @2.04± 0.38 and 4.04± 0.66 aphids plant<sup>-1</sup> in ZBNF and CF systems, during the last week of October, 2019 (44<sup>th</sup> SW), and peaked to 18.88± 3.94 and 28.38± 4.94 aphids plant<sup>-1</sup> during 6<sup>th</sup> and 5<sup>th</sup> SW in ZBNF and CF systems, respectively (Fig.1A). The

Table 1. Diversity indices of insect pests and their natural enemies in cauliflower under ZBNF and CF systems (2018-19, 2019-20)

Diversity indices	Insect-pests				Natural enemies			
	2018-19		2019-20		2018-19		2019-20	
	Farming systems	Farming systems	Farming systems	Farming systems	Farming systems	Farming systems	Farming systems	
	ZBNF	CF	ZBNF	CF	ZBNF	CF	ZBNF	CF
Simpson index	0.78	0.68	0.74	0.64	0.51	0.54	0.47	0.50
Shannon index (H)	0.73	0.99	0.82	1.06	1.23	0.78	1.38	0.83
H <sub>max</sub>	1.79	1.79	1.79	1.94	1.38	1.10	1.60	1.10
Evenness (J)	0.41	0.55	0.46	0.55	0.89	0.71	0.86	0.75
Dominance (D)	0.59	0.45	0.54	0.45	0.11	0.29	0.14	0.25

flea beetle appeared during the last week of October, 2018 (44<sup>th</sup> standard week) @  $0.14 \pm 0.05$  and  $0.30 \pm 0.07$  beetles plant<sup>-1</sup> in ZBNF and CF systems, and reached peak of  $0.28 \pm 0.08$  and  $0.60 \pm 0.12$  beetle plant<sup>-1</sup> during the 50<sup>th</sup> SW in ZBNF and CF systems, respectively; during the second year, the incidence appeared during the last week of October, 2019 (44<sup>th</sup> SW) @  $0.10 \pm 0.04$  and  $0.26 \pm 0.06$  beetle plant<sup>-1</sup> in ZBNF and CF systems respectively; these reached the peak in the second week of December, 2019 (50<sup>th</sup> SW) @  $0.24 \pm 0.07$  and  $0.58 \pm 0.13$  beetles plant<sup>-1</sup> in ZBNF and CF systems, respectively (Fig.1B).

Whitefly infestation started on 47<sup>th</sup> SW in ZBNF and CF systems, with the peak reaching in 52<sup>nd</sup> SW ( $0.58 \pm 0.16$  and  $1.28 \pm 0.28$  whiteflies plant<sup>-1</sup>) in ZBNF and CF systems, respectively in 2018-19; in 2019-20, the incidence appeared during the third week of November, 2019 (47<sup>th</sup> SW) and reached the peak in the last week of December, 2019 (52<sup>th</sup> SW) (Fig.1C). During 2018-19, Diamond back moth (DBM) started on 1<sup>st</sup> SW in ZBNF and CF systems, and reached peak in the third week of February, 2019 (8<sup>th</sup> SW) @  $0.96 \pm 0.16$  larvae plant<sup>-1</sup> in ZBNF system, and  $1.40 \pm 0.18$  larvae plant<sup>-1</sup> in CF system; during 2<sup>nd</sup> year, the incidence started during




Fig. 1. Seasonal abundance of insect-pests of cauliflower under ZBNF and CF systems during 2018-19 and 2019-20

the first week of January, 2020 (1<sup>st</sup> SW) and reached the peak of  $0.54 \pm 0.10$  larvae plant<sup>-1</sup> in ZBNF system, during the third week of February, 2020 (8<sup>th</sup> SW); while, in CF system, peak was at  $0.88 \pm 0.16$  larvae plant<sup>-1</sup> in the second week of February, 2020 (7<sup>th</sup> SW) (Fig. 1D). Likewise, cabbage head borer started infesting the crop during 3<sup>rd</sup> SW in both the systems in both the seasons with peak incidence being during the 8<sup>th</sup> SW (Fig. 1E). Cabbage butterfly infestation started in the first week of February, 2019 and 2020 i.e. 6<sup>th</sup> SW, with its larval counts reaching the peak in the fourth week of February, 2019 and 2020 (9<sup>th</sup> SW-  $6.52 \pm 2.37$  and  $16.00 \pm 3.97$  larvae plant<sup>-1</sup> in ZBNF and CF systems, 2019; and  $6.86 \pm 2.15$  and  $16.06 \pm 3.94$  larvae plant<sup>-1</sup> in 2020, respectively (Fig. 1F). Cabbage semilooper was recorded only in CF system and infestation started on 7<sup>th</sup> SW, with peak incidence recorded on 9<sup>th</sup> SW during second year (Fig. 1G). In case of natural enemies *D. rapae* parasitization was observed on 44<sup>th</sup> and 46<sup>th</sup> standard weeks in of ZBNF and CF systems during 2018-19 and its peak incidence ( $1.28 \pm 0.35$  and  $1.94 \pm 0.39$  mummified aphids plant<sup>-1</sup>) were recorded on 5<sup>th</sup> and 6<sup>th</sup> SW in CF and ZBNF systems. During 2019-20 the peak incidence of  $1.60 \pm 0.33$  and  $2.26 \pm 0.47$  mummified aphids plant<sup>-1</sup> was observed on 5<sup>th</sup> and 6<sup>th</sup> SW in CF and ZBNF systems, respectively.

The maximum parasitization (13.56 and 4.61%) was observed on 1<sup>st</sup> and 5<sup>th</sup> SW, 2019 in ZBNF and CF systems, respectively. During 2019-20 peak parasitization (22.22 and 6.40%) was observed on 52<sup>nd</sup> and 46<sup>th</sup> SW in ZBNF and CF systems, respectively (Fig. 2A). Activity of coccinellids started ( $0.10 \pm 0.04$  and  $0.08 \pm 0.04$  beetles plant<sup>-1</sup>) on 46<sup>th</sup> and 47<sup>th</sup> SW during 2018-19, and  $0.16 \pm 0.06$  and  $0.08 \pm 0.04$  beetles plant<sup>-1</sup> during 2019-20, in ZBNF and CF systems. Peak occurrence of coccinellids ( $0.58 \pm 0.12$  and  $0.28 \pm 0.09$  beetles plant<sup>-1</sup>) was observed during 6<sup>th</sup> and 7<sup>th</sup> SW in 2018-19, and ( $0.86 \pm 0.17$  and  $0.42 \pm 0.11$  beetles plant<sup>-1</sup>) during 2019-20, in ZBNF and CF systems (Fig. 2B). Syrphids appeared during 50<sup>th</sup> and 51<sup>st</sup> SW in ZBNF and CF systems during both seasons with maximum counts observed on 8<sup>th</sup> SW (Fig. 2C); *D. semiclausam* appeared on 5<sup>th</sup> SW in ZBNF, while it was not observed in CF system during both seasons- maximum was on 9<sup>th</sup> SW in ZBNF system in both seasons (Fig. 2D). Similarly, parasitic wasp, *C. glomerata* was seen only in ZBNF system during 2019-20 and this occurred during 8<sup>th</sup> SW with peak being in 9<sup>th</sup> standard week (Fig. 2E).

In ZBNF system organic amendment and intercropping had an impact on natural enemies and use of intercrop also repelled certain insect pests up to

a remarkable level. Sharma et al. (2020) also observed similar insect pests in their study from cauliflower, revealing 14 species of insect pests, of which seven are similar as in this study. Rana (2019) also reported these pests from the cruciferous ecosystem from Himachal Pradesh and revealed a better impact of ZBNF than CF system. Current study recovered all insect pests reported by Rana (2019) except green peach aphid *Myzus persicae* (Sulzer). Similar results were obtained by Mishra et al. (2018) where cabbage aphids (*B. brassicae*) appeared from 35<sup>th</sup> to 47<sup>th</sup> SW in cauliflower ecosystem. Mandal and Patnaik (2008) also observed insects of cole crops with peak incidence of *B. brassicae* during January and February. Patra et al. (2013) documented the seasonal abundance of DBM during late February to early March, which is in accordance with the present results. Singh et al. (2015) reported DBM from the 2<sup>nd</sup> SW and peak in 8<sup>th</sup> SW. Embaby (2015) observed similar incidence of *B. brassicae* and DBM. Dewanda and Khan (2016) also found DBM on cauliflower during the first week of September which gradually increased and reached its peak during first week of November. Similar results were reported by Meena and Singh (2012) on DBM that it appeared 35 days after transplanting of cabbage with peak in the last week of January. Das et al. (2017) studied the effect of temperature (°C) on species richness of whitefly in mustard- incidence was very low at 20 November, 2015, which gradually reached the peak in 19 March, 2016.

Khan and Talukder (2017) studied the influence of weather factors on the abundance and population dynamics of *P. brassicae* on cabbage- and found that larval incidence ranged from 0.58 to 1.98 larvae/ plant with peak on 5<sup>th</sup> February. Sharma et al. (2020) found *C. septempunctata* was the most abundant predator of aphids in cauliflower with relative proportion of 74.55 and 52.98% and the parasitization of *P. xylostella* by *D. semiclausam* was 28.94 and 30.77%; they also found that *C. glomerata* was the most abundant parasitoid of the larva of *P. brassicae* and *P. rapae* (42.43 and 57.58% during 2017 and 2018, respectively) which is quite similar to present results; and 10 predators and 4 parasitoids were observed in contrast to 5 predators and 3 parasitoids in the present study. The incidence of *P. brassicae* observed now is similar to that reported by Dwivedi et al. (2015). Jakhar and Singh (2018) observed the occurrence of coccinellids in cauliflower with *C. septempunctata* as the most dominant. Gaikwad et al. (2018) studied the insect pests and natural enemies of cauliflower during 2017 and reported the incidence of *B. brassicae*, *P. xylostella* and *H. undalis*. In addition, they






Fig.2. Seasonal abundance of natural enemies of insect pests of cauliflower under ZBNF and CF systems (2018-19, 2019-20)

recorded the occurrence of syrphid flies and mummified aphids. Evidences and results of current study provide a platform to conclude that ZBNF is a better farming system than CF system as it not only repels the pests, but conserves the natural enemies and improves the crop ecosystem.

#### ACKNOWLEDGEMENTS

The authors thank the Department of Agriculture, Government of Himachal Pradesh for funding the project on Subhash Palekar Natural Farming System under Prakritik Kheti Khushhal Yojna and the Professor and Head, Department of Entomology, Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan for providing facilities.

#### REFERENCES

- Chatterjee S S, Bose T K, Som M G. 1986. Cole crops. Vegetable crops in India. Naya Prakash. Calcutta. 165-247 pp.
- Das S, Rahman M, Kamal M, Shishir A. 2017. Species richness of thrips and whiteflies and their predators in mustard fields. Journal of the Bangladesh Agricultural University 15(1): 7-14.
- Dewanda P, Khan S. 2016. A field study of population dynamics of major insect pests and their natural enemies on cauliflower of Ajmer district. International Journal of Agriculture Science 8: 2642-2645.
- Dwivedi R K, Awaneesh C, Pal R K, Singh D K, Akhilesh T. 2015. Seasonal abundance of cabbage (*Brassica oleracea* var. capitata L.) butterfly (*Pieris brassicae*) and its management. Current Advances in Agricultural Sciences (An International Journal) 4(1): 87-89.
- Embaby E S M, Lotfy D E S. 2015. Ecological studies on cabbage pests. International Journal of Agricultural Technology 11(5): 1145-1160.
- Gaikwad A D, Bhede B V, Bokan S C, Bhosle B B. 2018. Seasonal incidence of major insect pests, natural enemies on cauliflower and

- their correlation with weather parameters. *Journal of Entomology and Zoology Studies* 6(5): 952-956.
- Gallo J, Pekár S. 1999. Winter wheat pests and their natural enemies under organic farming system in Slovakia: Effect of ploughing and previous crop. *Anz Schadlingskde Pflanzenschutz Umweltschutz* 72: 31-36.
- Jakhar M, Singh S K. 2018. Study of seasonal abundance of diamondback moth and natural enemies in cauliflower, crop and their relation with the environmental factors. *Journal of Entomology and Zoology Studies* 6(6): 1250-1252.
- Khan M M H, Talukder S. 2017. Influence of weather factors on the abundance and population dynamics of *Spodoptera litura* F. and *Pieris brassicae* L. on cabbage. *SAARC Journal of Agriculture* 15(1): 13-21.
- Liu S S, Wang X G, Guo S J, He J H, Shi J H. 2000. Seasonal abundance of the parasitoid complex associated with the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae) in Hangzhou, China. *Bulletin of Entomological Research* 90: 221-231.
- Mandal S M A, Patnaik N C. 2008. Interspecific abundance and seasonal incidence of aphids and aphidophagous predators associated with cabbage. *Journal of Biological Control* 22(1): 195-198.
- Meena B L, Dadhich S R, Kumawat R L. 2002. Efficacy of some insecticides against ladybird beetle, *Coccinella septumpunctata* L. feeding on fenugreek aphid, *Acyrtosiphon pisum* (Harris). *Ann Biol* 18: 171-173.
- Meena S C, Singh V. 2012. Seasonal incidence of *Plutella xylostella* L. on cabbage in arid region of Rajasthan. *Annals of Plant Protection Sciences* 20(2): 326-328.
- Mishra R, Mhaske B M, Patil N M, Desale S B. 2018. Influence of abiotic factors on the seasonal incidence of major insect pests on cauliflower. *Journal of Entomology and Zoology Studies* 6(5): 2085-2088.
- Palekar S. 2016. Vegetable crops in zero budget spiritual farming. Zero budget spiritual agriculture system research, development and extension movement, Amaravati, Maharashtra.
- Patra S, Dhote V W, Alam S K F, Das B C, Chatterjee M L, Samanta A. 2013. Population dynamics of major insect pests and their natural enemies on cabbage under new alluvial zone of West Bengal. *The Journal of Plant Protection Sciences* 5(1): 42- 49.
- Raja M, William John, David B V. 2014. Population dynamics of key insect pests of cabbage in Tamil Nadu. *Indian Journal of Entomology* 76(1): 01-07.
- Rana A. 2019. Insect-pests, natural enemies and soil micro-flora in cabbage under zero budget natural farming and conventional farming systems. M Sc Thesis. Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan.
- Shannon C E. 1948. A mathematical theory of communication. *Bell System Technical Journal* 27: 379-423.
- Sharma S, Verma S C, Sharma P L, Chandel R S. 2020. Diversity of inset-pests and their natural enemies in cauliflower under mid hills of Himachal Pradesh. *Journal of Entomology and Zoology Studies* 8(2): 1204-1209.
- Sheoran O P, Tonk D S, Kaushik L S, Hasija R C, Pannu R S. 1998. Statistical software package for agricultural research workers.
- Simpson E H. 1949. Measurement of diversity. *Nature* 163: 688.
- Singh R P, Mishra M K, Chandra U, Pratap N. 2015. A field study of pest of cauliflower in eastern plain zone areas. *Plant Archives* 15(1): 561-564.
- Yadav D P. 1989. Integrated pest management on mustard. *Annals of Agricultural Research* 22: 429-431.

(Manuscript Received: February, 2021; Revised: March, 2021;

Accepted: July, 2021; Online Published: October, 2021)

Online published (Preview) in [www.entosocindia.org](http://www.entosocindia.org) Ref. No. e21031