



EFFICACY OF MINERAL AND NON-EDIBLE SEED OILS AGAINST APHIDS AND WHITEFLY IN POTATO

MURLIDHAR SADAWARTI*, SUBHASH KATARE, S P SINGH, M ABAS SHAH¹,
R K SAMADHIYA, SANJEEV SHARMA² AND R K SINGH²

ICAR-Central Potato Research Institute-Regional Station, Gwalior 474020, Madhya Pradesh, India

¹ICAR-Central Potato Research Institute-Regional Station, Jalandhar 144003, Punjab, India

²ICAR-Central Potato Research Institute, Shimla 171001, Himachal Pradesh, India

*Email: murlidharsada@gmail.com (corresponding author)

ABSTRACT

Field experiments were conducted at the ICAR-Central Potato Research Institute-Regional Station, Gwalior, Madhya Pradesh during 2015-16 and 2016-17 to evaluate efficacy of mineral and non-edible seed oils against green-peach aphid *Myzus persicae* and cotton whitefly *Bemisia tabaci* in potato. Three sprays were done at 15 days interval revealed that maximum reduction in their incidence was observed with imidacloprid 17.8SL (0.03%) followed by mineral oil (6 ml/l); *B. tabaci* incidence reduced by 72.63% and 61.06%, respectively, while it was 87.60% and 62.27% with *M. persicae*. Similar trend was observed with all the three sprays. The non-edible oils were not effective. Incidence of viruses was the least (0.26 and 0.27%, respectively) with imidacloprid 17.8SL (0.03%) and mineral oil (6 ml/l). Thus, imidacloprid 17.8SL (0.03%) and mineral oil (6 ml/l) can be recommended to manage vector-virus complex in potato.

Key words: *Myzus persicae*, *Bemisia tabaci*, potato, Kufri Jyoti, non-edible oil, mineral oil, imidacloprid, sprays, potato virus, vector virus complex, yield

India is the second largest producer of potato in the world (FAOSTAT, 2017), with >85% of being grown in the vast Indo-Gangetic plains of north India (subtropics) during short winter days from October to March (Khurana and Naik 2003). Potato is infested by a number of aphids transmitting more than three dozen viruses, affecting the yield and quality (Salazar, 2006; Bhatnagar et al., 2017). Peach-potato aphid *Myzus persicae* (Sulzer) is the most efficient aphid vector in potato (Bhatnagar et al., 2012). In addition, the cotton whitefly *Bemisia tabaci* (Gennadius) is a major constraint for healthy seed potato production as it transmits the Tomato leaf curl New Delhi virus (potato) which results huge yield loss and degeneration of seed stocks (Shah et al., 2019). Currently, the management of vector-virus complex in potato depends on synthetic pesticides, in addition to the cultural practice of growing the crops during low aphid activity period (Pushkarnath, 1959; 1967). However, the management of non-persistent viruses like potato virus Y (PVY) continues to be a challenge. Mineral oils are known as effective means to control insect pests like aphids and to reduce the spread of non-persistent viruses (Shah et al., 2021; Perring et al., 1999). These are widely used in Europe and are increasingly used in the United States and Eastern Canada for seed potato production.

Several studies have shown that mineral oils interfere with virus retention in the aphid mouthparts (stylet) and foliar application of mineral oil reduced the transmission of PVY to potatoes (Boiteau et al., 2008; Najar-Rodriguez et al., 2008; Wrobel, 2009). In addition to this, various plant-derived oils have been shown to be effective against the vectors e.g., neem oil (Isman, 2006); pongamia/ karanj oil (Kumar and Singh, 2002). There is little knowledge about the effect of mineral, neem and karanj oils on the sucking pests of potato under north-central Indian conditions, and hence the present study.

MATERIALS AND METHODS

Field experiments were conducted at the ICAR-Central Potato Research Station, Gwalior, during 2015-16 and 2016-17 on potato cv. Kufri Jyoti. The experiments were laid in randomized complete block design, with six treatments and four replications. The treatments included mineral oil (3 ml/l and 6 ml/l), pongamia oil (6 ml/l), neem oil (7.3 ml/l), imidacloprid 17.8SL (3 ml/10l) as standard check along with untreated control. Total of three sprays were given at 15 days interval, starting from 40 days. The crops were raised with recommended agronomic practices, with crop planted on 09/11/2015 and 25/10/2016 during the

two seasons. Observations on the number of adult *B. tabaci*, and adults and 3rd and 4th instar nymphs of *M. persicae* were taken from ten randomly selected plants/plot, one day before, and 7 and 14 days after treatment (DAT). The numbers were counted from three leaves on each plant, one each from upper, middle and lower strata, preferably in the early morning hours. The reduction in incidence was calculated as per Henderson and Tilton (1955). The total tuber yield was recorded on whole plot basis, and % incidence of viruses (severe and mild mosaic, leaf roll) was taken from visual observation on all the plants in each plot. The data on number of insects, % reduction in incidence over control and tuber yield were subjected to ANOVA after appropriate transformation. The treatment means were separated by least significant difference (LSD) test.

RESULT AND DISCUSSION

Observation on *B. tabaci* revealed that the pretreatment counts ranged from 4.04 to 5.19/ 10 plants which increased till mid-December and declined sharply afterwards, again appearing towards February. Therefore, three spray treatments were given; two in November-December and one in February; and maximum reduction in incidence was observed with imidacloprid 17.8 SL (0.03%) followed by that of mineral oil (6 ml/ l). Similar consistent trend was noted for all the three sprays- its incidence reduced by 72.63% and 61.06%, respectively with these treatments. The non-edible oils did not cause an appreciable reduction, while treatments with pongamia oil gave a reduction of 51.20% and neem oil s 55.16% reduction (Table 1). These results are in conformity with those

Table 1. Efficacy of mineral and non-edible seed oils against *B. tabaci* and *M. persicae* in potato

Treatments	Mean no. of from 10 plants (3 leaves/ plant)									% reduction over control
	1 st Spray			2 nd Spray			3 rd Spray			
	Pre-count	7 DAT	14 DAT	Pre-count	7 DAT	14 DAT	Pre-count	7 DAT	14 DAT	
<i>B. tabaci</i>										
Control	4.19 (2.27)	8.98 (3.15)	10.30 (3.35)	10.30 (3.35)	6.10 (3.35)	3.80 (2.18)	3.80 (2.18)	3.25 (1.98)	5.33 (2.51)	-
Imidacloprid 17.8SL @ 0.03%	4.04 (2.24)	3.00 (1.89)	1.75 (1.61)	1.75 (1.61)	0.00 (1.00)	0.38 (1.14)	0.38 (1.14)	0.13 (1.05)	0.17 (1.07)	72.63
Mineral oil @ 3 ml/ l	4.57 (2.35)	7.75 (2.95)	4.25 (2.21)	4.25 (2.21)	1.75 (1.65)	0.50 (1.20)	0.50 (1.20)	0.25 (1.10)	0.42 (1.17)	42.97
Mineral oil @ 6 ml/ l	4.64 (2.37)	5.25 (2.45)	1.88 (1.61)	1.88 (1.61)	0.54 (1.23)	0.17 (1.07)	0.17 (1.07)	0.00 (1.00)	0.17 (1.07)	61.06
Pongamia oil @ 6.8 ml/ l	4.94 (2.43)	5.88 (2.59)	5.50 (2.52)	5.50 (2.52)	2.25 (1.77)	0.50 (1.20)	0.50 (1.20)	0.00 (1.00)	0.50 (1.20)	51.20
Neem oil @ 7.3 ml/ l	5.19 (2.48)	6.88 (2.78)	4.13 (2.24)	4.13 (2.24)	1.25 (1.49)	0.50 (1.20)	0.50 (1.20)	0.25 (1.10)	0.25 (1.10)	55.16
SEm	0.002	0.23	0.22	0.22	0.10	0.11	0.11	0.13	0.09	
CD (p = 0.05)	NS	0.71	0.68	0.68	0.30	0.35	0.35	0.42	0.27	
<i>M. persicae</i>										
Control	1.75 (1.65)	5.13 (2.47)	10.50 (3.38)	10.50 (3.38)	12.68 (3.69)	14.38 (3.69)	14.38 (3.69)	15.00 (3.99)	12.75 (3.68)	-
Imidacloprid 17.8SL @ 0.03%	2.16 (1.77)	2.43 (1.84)	1.00 (1.36)	1.00 (1.36)	0.25 (1.10)	0.25 (1.10)	0.25 (1.10)	0.00 (1.00)	0.00 (1.00)	87.60
Mineral oil @ 3 ml/ l	1.90 (1.70)	4.13 (2.26)	5.50 (2.54)	5.50 (2.54)	3.13 (2.00)	4.63 (2.00)	4.63 (2.00)	3.25 (2.05)	1.75 (1.65)	44.19
Mineral oil @ 6 ml/ l	2.28 (1.81)	3.25 (2.05)	4.13 (2.26)	4.13 (2.26)	1.75 (1.65)	2.38 (1.65)	2.38 (1.65)	1.50 (1.57)	0.25 (1.10)	62.27
Pongamia oil @ 6.8 ml/ l	1.98 (1.72)	2.13 (1.60)	6.75 (2.76)	6.75 (2.76)	4.75 (2.39)	3.75 (2.39)	3.75 (2.39)	1.75 (1.64)	2.00 (1.73)	49.66
Neem oil @ 7.3 ml/ l	2.09 (1.75)	4.50 (2.23)	4.38 (2.64)	4.38 (2.64)	3.00 (1.99)	2.75 (1.99)	2.75 (1.99)	1.75 (1.65)	0.50 (1.20)	51.58
SEm	0.038	0.23	0.13	0.13	0.10	0.10	0.10	0.08	0.13	
CD (p = 0.05)	NS	NS	0.40	0.40	0.30	0.30	0.30	0.24	0.40	

Values pooled data of two seasons; Precount- Pretreatment count; DAT – Days after treatment; values in parentheses square root transformed as $\sqrt{x+0.5}$; NS = not significant

of Bhatnagar et al. (2016). Imidacloprid 17.8SL (150 ml/ ha) at 15 days interval was the most effective (Nag et al., 2018). About the *M. persicae*, it appeared after mid-December, therefore, three sprays were made from December to February, which increased from 1.75/ 10 plants in the third week of December to 15.00/ 10 plants towards mid-February and declined afterwards. The data on the incidence pooled over two seasons revealed that maximum reduction was obtained with imidacloprid 17.8SL followed by that of mineral oil (6 ml/ l), with results being similar with all the three sprays, with incidence reducing by 87.60% and 62.27%, respectively. The non-edible oils did not show such reduction (Table 1). These results are in confirmation with those of Khan (2011), Basavaraju et al.(2015) and Nag et al. (2018). Azadirachtin results in >60% suppression of aphids (Ghosh, 2015). Neem oil (1%), NSKE (5%), pongamia (1-2%) and karanj oil (1 and 2%) are reported as effective (Kumar and Singh, 2002; Pavela, 2007). Although the incidence of viruses was low during both the seasons, significant reduction in the incidence being with treated plots, the least incidence being with imidacloprid 17.8 SL (0.03%) and mineral oil (6 ml/ l). Total tuber yield (ranging 29.22 to 31.44 mt/ ha) was non significant among the treatments.

REFERENCES

- Basavaraju B S, Chakravarthy A K, Siddagangaiah, K N. 2015. Evaluation of insecticides against potato aphid, *Myzus persicae* Sulzer and thrips, *Thrips palmi* Karney. Karnataka Journal of Agricultural Science 28(4): 524-527.
- Bhatnagar A, Somani A K, Chander S. 2012. Buildup of *Myzus persicae* and development of mosaics on potato in central India. Potato Journal 39(2): 208-210.
- Bhatnagar A. 2016. Effectiveness of non-edible oils on insect vector and disease incidence in potato crop- ITK technique. Arora S, J P Sharma, S Chakravorty, N Sharma, P Joshi (eds.). Indigenous technologies in plant protection. ICAR- National Research Centre for Integrated Pest Management, New Delhi. pp. 75-82.
- Bhatnagar A, Pant R P, Sridhar J, Chakrabarti S K, Lal M. 2017. Incidence of apical leaf curl disease (ToLCNDV), and economics and reaction of potato (*Solanum tuberosum*) cultivars against whitefly, *Bemisia tabaci* in northern India. Indian Journal of Agricultural Science 87(12): 1673-1678.
- Boiteau G, Singh M, Lavoie J. 2008. Crop border and mineral oil sprays used in combination as physical control methods of the aphid-transmitted potato virus Y in potato. Pest Management Science 65: 255-259.
- FAOSTAT. 2017. Statistical database. <http://faostat.fao.org/>
- Ghosh S K. 2015. Integrated field management of aphids (*Myzus persicae* Sulz. And *Aphis gossypii* Glov. together) on potato (*Solanum tuberosum* L.) using bio-pesticides. International Journal of Science, Environment and Technology 4(3): 682-689.
- Henderson C F, Tilton E W. 1955. Tests with acaricides against the brown wheat mite. Journal of Economic Entomology 48: 157-161.
- Isman MB. 2006. Botanical insecticides, deterrents and repellants in modern agriculture and increasing regulated world. Annual Review of Entomology 51: 45-66.
- Khan MA, Saljoqi A U R, Hussain N, Sattar S. 2011. Response of *Myzus persicae* (Sulzer) to imidacloprid and thiamethoxam on susceptible and resistant potato varieties. Sarhad Journal of Agriculture 27(2): 263-270.
- Khurana S M P, Naik P S. 2003. The potato: an overview. pp. 1-14. Khurana S M P, J S Minhas, S K Pandey (eds.). The potato production and utilization in subtropics. Mehta Publishers, New Delhi.
- Kumar M, Singh R. 2002. Potential of *Pongamia glabra* Vent as an insecticide of plant origin. Biological Agriculture and Horticulture 20: 29-50.
- Nag D, Gupta R, Bhargav P, Kumar Y, Bisen M S. 2018. Relative efficacy of insecticides against potato aphid (*Myzus persicae* Sulzer), whitefly (*Bemisia tabaci* Genn.) and ladybird beetle. International Journal of Chemical Studies 6(1): 1182-1186.
- Najar-Rodriguez, A J, Lavidis N A, Mensah R K, Choy P T, Walter G H. 2008. The toxicological effects of petroleum spray oils on insects-evidence for an alternative mode of action and possible new control options. Food and Chemical Toxicology 46: 3003-3014.
- Pavela R. 2007 Possibilities of botanical insecticide exploitation in plant protection. Pest Technology 1(1): 47-52
- Perring T M, Gruenhagen N M, Farrar C A. 1999. Management of plant viral diseases through chemical control of insect vectors. Annual Review of Entomology 44: 457-481.
- Pushkarnath 1959. Producing healthy seed potatoes in the plains: A new approach. Indian Potato Journal 1: 63-72.
- Pushkarnath 1967. Seed potato production in the sub-tropical plains of India. American Potato Journal 44: 429-441.
- Salazar L F. 2006. Virus diseases of potatoes. Encyclopedia of life sciences. Wiley, Chichester. <https://doi.org/10.1038/ngp.els.0004300>.
- Shah MA, Malik K, Bhatnagar A, Katare S, Sharma S, Chakrabarti SK. 2019. Effect of temperature and cropping sequence on the infestation pattern of *Bemisia tabaci* in potato. Indian Journal of Agricultural Science 89(11): 1802-1807.
- Shah M A, Naga K C, Subhash S, Sharma S, Kumar R. 2021. Use of petroleum-derived spray oils for the management of vector-virus complex in potato. Potato Research. <https://doi.org/10.1007/s11540-021-09505-0>
- Wrobel S. 2009. The retention of PVY in the stylet of *Myzus persicae* Sulz. after the application of mineral oil on potato plants. Plant Breeding and Seed Science 60: 3-12.