



EFFICACY OF SULFOXAFLOR 12%SC AGAINST APHIDS COMPLEX AND *COCCINELLA SEPTEMPUNCTATA* L. IN WHEAT

SUBHASH KATARE, POONAM JASROTIA*, JAYANT YADAV¹ AND M S SAHARAN

ICAR-Indian Institute of Wheat and Barley Research, Karnal 132001, Haryana, India

¹Department of Entomology, C.C.S. Haryana Agricultural University, Hisar 125004, Haryana, India

*Email: poonamjasrotia@gmail.com (corresponding author)

ABSTRACT

Efficacy of dosages of sulfoxaflor 12%SC were evaluated for the control of foliar aphids under Contract Research Project on wheat during rabi seasons of 2014-15 and 2015-16 at the research farm of ICAR-Indian Institute of Wheat and Barley Research, Karnal, Haryana. The standard check insecticides thiamethoxam 25WDG and quinalphos 25EC were included along with sulfoxaflor, and all the treatments were found to be significantly superior. The incidence of aphids/ shoot/ plant was significantly less with sulfoxaflor 12 SC @ 24, 27 and 30 g a.i./ ha with 94.54, 95.27 and 96.03% reduction, respectively; these being at par with each other, followed by thiamethoxam 25WDG @ 12.5 g a.i./ ha and quinalphos 25EC @ 250 g a.i./ ha. The seven-spotted ladybird beetle *Coccinella septempunctata* L. was the main natural enemy observed in the crop, and its counts were the least (1.50/ m²) with thiamethoxam 25WDG @ 12.5 g a.i./ha at 14 DAS. The pooled data revealed that the yield was significantly more with sulfoxaflor 12SC @ 24, 27 and 30 g a.i./ ha (52.21, 52.62 and 54.32 q/ ha) followed by thiamethoxam 25WDG @ 12.5 g a.i./ ha (49.87 q/ ha) which was at par with quinalphos 25EC @ 250 g a.i./ ha (47.73 q/ha). Considering incremental cost benefit ratio, sulfoxaflor 12% SC @30 g a.i/ ha (1:2.70) is the most economical, and no phytotoxicity symptoms were observed at its doses @30 and 60 g a.i/ ha.

Key words: Insecticides, sulfoxaflor, thiamethoxam, quinalphos, wheat, *Rhopalosiphum maidis*, *Rhopalosiphum padi*, coccinellids, *Coccinella septempunctata*, phytotoxicity, yield, cost benefits

Wheat (*Triticum aestivum* L.) is one of the most important cereal crops and the staple food throughout the world. Among the different insect pest attacking wheat, cereal aphids are the most widely distributed and are posing a serious threat (Singh, 1983; Yadav, 2003). The major aphid species found in wheat are *Rhopalosiphum maidis* (Fitch), *Rhopalosiphum padi* L., *Sitobion miscanthi* and *Sitobion avenae* (F.) (Jasrotia et al., 2021; Singh et al., 2020) They cause direct damage by sucking cell sap of leaves, young shoots, causing distortion, stunting, leaf curling, wilting and twisting and also cause indirect damage by secreting honey dew on leaves which reduce photosynthetic activity and attract sooty mould growth and premature leaf senescence (Deol et al., 1987; Ozder, 2002;). The aphids caused 10 to 50% reduction in yield. The damage is severe in cold and cloudy weather during winters. They appear mostly from December to January (Trdan and Milevoj; 1999; Singh and Deol, 2003; Katare et al., 2018). Management of aphids has been done primarily with chemical methods. Wide spread use of the pesticides causes several socio-economic problems. More than 20 aphid species have showed resistance to carbamate, pyrethroid and organophosphate based insecticides (Yu

et al., 2016). New insecticide formulations with high efficacy and novel mode of action are being used to control aphids in wheat crop (Longhurst et al., 2013; Bhanu et al., 2015; Katare et al., 2015). New chemicals such as flubendamide 480SC, thiamethoxam 25WG and chlorantranilipride 18.5SC, and entomopathogenic fungi, *Metarhizium anisopliae*, *Verticillium lecanii* have been tried against aphids in wheat and barley and have been found effective (Katare et al., 2017; 2018; Jasrotia et al., 2018). However, scanty information is available on the efficacy of newer insecticide, sulfoxaflor 12%SC against wheat aphids, and the present study evaluates its efficacy against wheat aphids.

MATERIALS AND METHODS

A field experiment was conducted at ICAR- Indian Institute of Wheat and Barley Research, Karnal (22.93°N, 88.53°E, 9.75 masl) during 2014-15 and 2015-16. Sulfoxaflor 12% SC w/v (11.4% SC w/w) was evaluated as foliar spray. The experiment was carried out in the 3rd week of November in randomized block design with eight treatments viz., T₁-Sulfoxaflor 12%SC (21 g a.i/ ha), T₂-Sulfoxaflor 12%SC (24 g a.i/ ha), T₃-Sulfoxaflor 12%SC (27 g a.i/ ha), T₄-Sulfoxaflor

12%SC (30 g a.i/ ha), T₅-Thiamethoxam 25%WDG (12.5 g a.i/ ha), T₆-Quinalphos 25%EC (250 g a.i/ ha), T₇-Sulfoxaflor 12%SC (60 g a.i/ ha) and T₈-Untreated control; and with three replications using variety HD 2967. A plot size of 7.5 m² was maintained, and standard package of practices followed (Kumar et al., 2001; Anonymous, 2014). The insecticidal sprays were applied just after mean infestation reached 20 aphids/ shoot/ plant or above. Five shoots from each treatment were selected randomly for recording the aphid incidence, with pre-count done 24 hr before spray and post-counts at 1, 3, 7 and 14 days after treatment (DAT). Only live aphids were counted, and of *R. maidis* and *R. padi*. The natural enemy that was observed was the seven-spotted ladybird beetle *Coccinella septempunctata* L. The occurrence of this beetle (adults and grubs) was observed in 1 m² quadrat in three spots in treated plots. At maturity, grain yield was recorded on whole plot basis and converted in q/ ha. Sulfoxaflor 12%SC (30g a.i/ha) (T₄) was evaluated for both efficacy and phytotoxicity while sulfoxaflor 12%SC (60g a.i/ ha) (T₇) for only phytotoxicity. For benefit cost analysis, the cost of insecticidal application was calculated as cost of insecticide+ labour charges+ machine cost etc. Similarly, the price of harvested wheat was calculated as per the prevailing market rate. The data were subjected to ANOVA and treatment means were separated by least significant difference test (Gomez and Gomez, 1984).

RESULTS AND DISCUSSION

The pooled data for 2014-15 and 2015-16 revealed that before spraying, incidence ranged from 33.20 to 30.20 aphids/ tiller/ plant and did not vary significantly. All the treatments were significantly superior over the control in respect of reducing foliar aphids. One day after the spray, least counts (4.00 aphids/ tiller/ plant) were observed in T₄ (sulfoxaflor 12%SC @ 30 g a.i/ ha). However, it was statistically at par with its dose @ 27g a.i/ ha- 4.43 aphids/ tiller/ plant and also @ 24 g a.i/ ha- 4.87 aphids/ tiller/ plant. Three days after the spray, again T₄ i.e was superior (0.87 aphids/ tiller/ plant) but statistically at par with its dose at 27 g a.i/ ha (1.30 aphids/ tiller/ plant) (Table 1). At 7 and 14 days after the spray, similar results were obtained followed by standard T₅ (thiamethoxam 25WDG @ 12.5 g a.i/ ha) and T₆ (quinalphos 25EC @ 250 g a.i/ ha). In untreated control plot, maximum number of aphids/ tillar/ plant were observed as 33.43, 33.43, 33.47 and 27.27 at 1, 3, 7 and 14 days after spray, respectively. The occurrence of *C. septempunctata* was non-significant at precount, and the least (1.50/ m²) in the standard check T₅ at 14 days after spraying (Table 2).

These observations agree with those of Chandi (2019). Comprehensive evaluation of field efficacy indicated that sulfoxaflor had quick acting properties and long persistent effects and it should be used

Table 1. Efficacy of doses of ssulfoxaflor 12% SC against foliar aphids in wheat

S. No.	Treatments	Dose (g a.i./ ha)	Before spray (BS) (Mean no. of aphids/ shoot/ plant)			1 Day after spray (Mean no. of aphids/ shoot/ plant)			3 Day after spray (Mean no. of aphids/ shoot/ plant)		
			2014-15	2015-16	Pooled mean	2014-15	2015-16	Pooled mean	2014-15	2015-16	Pooled mean
T ₁	Sulfoxaflor 12% SC	21	30.20 (5.54)*	30.33 (5.54)	30.27 (5.54)	8.40 (2.98)	2.73 (1.80)	5.57 (2.46)	3.07 (1.89)	1.67 (1.47)	2.37 (1.69)
T ₂	Sulfoxaflor 12% SC	24	31.60 (5.66)	27.00 (5.24)	29.30 (5.46)	7.80 (2.88)	1.93 (1.55)	4.87 (2.32)	2.40 (1.70)	1.20 (1.30)	1.80 (1.52)
T ₃	Sulfoxaflor 12% SC	27	32.53 (5.75)	31.33 (5.63)	31.93 (5.69)	7.00 (2.74)	1.87 (1.53)	4.43 (2.22)	1.60 (1.45)	1.00 (1.22)	1.30 (1.34)
T ₄	Sulfoxaflor 12% SC	30	29.40 (5.45)	28.27 (5.36)	28.83 (5.41)	6.80 (2.70)	1.20 (1.30)	4.00 (2.12)	0.93 (1.19)	0.80 (1.14)	0.87 (1.17)
T ₅	Thiamethoxam 25% WDG	12.5	32.07 (5.70)	31.33 (5.64)	31.70 (5.67)	8.40 (2.98)	2.20 (1.64)	5.30 (2.41)	3.93 (2.10)	1.47 (1.39)	2.70 (1.79)
T ₆	Quinalphos 25% EC	250	32.80 (5.76)	28.27 (5.36)	30.53 (5.57)	11.60 (3.47)	4.20 (2.16)	7.90 (2.90)	7.67 (2.86)	1.87 (1.54)	4.77 (2.29)
T ₇	Untreated Check	-	33.20 (5.80)	29.27 (5.45)	31.23 (5.63)	35.30 (5.97)	31.67 (5.66)	33.43 (5.82)	38.20 (6.22)	28.67 (5.40)	33.43 (5.82)
	SE ±		0.14	0.16	0.11	0.09	0.12	0.07	0.08	0.08	0.07
	CD (p=0.05)		NS	NS	NS	0.27	0.36	0.22	0.25	0.25	0.20

* Figures in parentheses $\sqrt{n+0.5}$ values

Table 2. Efficacy of doses of Sulfoxaflor 12%SC against foliar aphids in wheat

Sr. No.	Treatments	Dose (g a.i./ha)	7 Days after spray (Mean no. of aphids/shoot/plant)			14 Days after spray (Mean no. of aphids/shoot/plant)			Pooled mean population (2014-15 to 2015-16)	% reduction over control	C. <i>septempunctata</i> /m ² (Pooled mean 2014-15 to 2015-16)	
			2014-15	2015-16	Pooled mean	2014-15	2015-16	Pooled mean			Before spray	14 Day After Spray
T ₁	Sulfoxaflor 12% SC	21	0.80 (1.14)*	1.13 (1.27)	0.97 (1.21)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	2.23	93.03	1.33 (1.35)	2.77 (1.81)
T ₂	Sulfoxaflor 12% SC	24	0.00 (0.71)	0.60 (1.05)	0.30 (0.89)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	1.74	94.54	1.51 (1.42)	2.14 (1.62)
T ₃	Sulfoxaflor 12% SC	27	0.00 (0.71)	0.60 (1.05)	0.30 (0.89)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	1.51	95.27	1.51 (1.42)	2.08 (1.61)
T ₄	Sulfoxaflor 12% SC	30	0.00 (0.71)	0.40 (0.95)	0.20 (0.84)	0.00 (0.71)	0.00 (0.71)	0.00 (0.71)	1.27	96.03	1.60 (1.45)	2.01 (1.58)
T ₅	Thiamethoxam 25% WDG	12.5	1.13 (1.28)	1.60 (1.44)	1.37 (1.36)	2.00 (1.58)	0.00 (0.71)	1.00 (1.22)	2.59	91.88	1.27 (1.33)	1.50 (1.41)
T ₆	Quinalphos 25% EC	250	7.53 (2.83)	2.40 (1.70)	4.97 (2.34)	4.33 (2.20)	0.00 (0.71)	2.17 (1.63)	4.95	84.48	1.45 (1.40)	1.57 (1.44)
T ₇	Untreated Check	-	40.50 (6.38)	26.73 (5.22)	33.47 (5.83)	35.20 (5.97)	19.33 (4.44)	27.27 (5.27)	31.90	-	1.73 (1.49)	5.49 (2.45)
	SE ±		0.06	0.07	0.03	0.05	0.07	0.03	-	-	0.04	0.04
	CD (p=0.05)		0.18	0.20	0.10	0.17	0.23	0.11	-	-	NS	0.12

*Figures in parentheses $\sqrt{n+0.5}$ values

Table 3. Economics of treatments (pooled data, 2014-15, 2015-16)

Sr. No.	Treatments	Dose (g a.i./ha)	Yield (q/ha)	Additional yield over control q/ha	Additional income over control (Rs.)	*Cost of cultivation (Rs.)	Gross income (Rs.)	Net income (Rs.)	B:C ratio
T ₁	Sulfoxaflor 12% SC	21	50.29	6.52	8802	26207	67892	41685	1: 2.59
T ₂	Sulfoxaflor 12% SC	24	52.21	8.44	11394	26467	70484	44017	1: 2.66
T ₃	Sulfoxaflor 12% SC	27	52.62	8.85	11947	26812	71037	44225	1: 2.65
T ₄	Sulfoxaflor 12% SC	30	54.32	10.55	14242	27187	73332	46145	1: 2.70
T ₅	Thiamethoxam 25% WDG	12.5	49.87	6.10	8235	26567	67325	40758	1: 2.53
T ₆	Quinalphos 25% EC	250	47.73	3.96	5346	26422	64436	38014	1: 2.44
T ₇	Untreated Check	-	43.77	-	-	23937	59090	35153	1: 2.47
	SE ±		1.26	-					
	CD (p=0.05)		3.88	-					

Total cost of cultivation except insecticide control: Rs. 23495/-; Rs. 24378/- (Avg. Rs. 23937/-); Price of wheat grains: Rs. 1200/q; Rs.500/- (Avg. Rs.1350/-); *Cost of cultivation = Cost of insecticide + labour + insecticide application + machine charges; Cost of insecticides: 1. Sulfoxaflor 12% SC: (Rs. 2270, 2530, 2875, 3250/ha) 2. Thiamethoxam 25% WDG: (Rs.2630/ha) 3. Quinalphos 25% EC: (Rs. 2485/ha)

alternatively with other insecticides for the control of wheat aphids (Babcock et al., 2011; Xin et al., 2019). Annetts et al. (2012) observed that sulfoxaflor demonstrated very good control of all major species of aphids infesting cereals, like oat aphid (*R. padi*), corn aphid (*R. maidis*), grain aphid (*S. miscanthi*) and rose grain aphid (*Metopolophium dirhodum*). The pooled data revealed that among the treatments % reduction of aphids ranged between 84.48 to 96.03%, with

maximum being with T₄ (96.03); and standard check T₅ (thiamethoxam 25WDG @ 12.5 g a.i./ ha) and T₆ (quinalphos 25EC @ 250 g a.i./ ha) led to 84.15 and 76.37% reduction, respectively (Table 2). The data for phytotoxicity were observed at 1, 7 and 14 DAT. The observations for the specific parameters like chlorosis, necrosis, wilting, hyponasty and epinasty were noted. There were no adverse effects of higher doses of sulfoxaflor 12% SC @ 30 and 60 g a.i/ ha on plants.

The pooled data showed that yield was significantly higher in treated plots, and significantly higher yield was obtained in T₄, T₃ and T₂ (54.32, 52.62 and 52.21 q/ha, respectively) and was at par with standard T₅ (49.87 q/ha) and T₆ (47.73 q/ha). Considering incremental cost benefit ratio (1:2.70) T₄ (sulfoxaflor 12%SC @ 30 g a.i/ ha) was found economical (Table 3). Therefore, present study revealed that being safer and economically cheapest, sulfoxaflor 12SC @ 30 g a.i./ ha can be used for the effective management of foliar aphids in wheat crop.

ACKNOWLEDGEMENTS

The authors thank the Director and PI, Division of Crop Protection, ICAR-IIWBR, Karnal, Haryana for providing necessary facilities.

REFERENCES

- Annetts R, Elias N, Corr I. 2012. Transform™ insecticide (sulfoxaflor) for control of aphids in canola. Proceedings. 16th ASA conference. Capturing opportunities and overcoming obstacles in Australian agronomy, Armidale, Australia. 1-4 pp.
- Anonymous 2014. Package of practices for crops of Punjab. Rabi 2014. Punjab Agricultural University, Ludhiana.
- Babcock J M, Gerwick C B, Huang J X, Loso M R, Nakamura G, Nolting S P, Rogers R B, Sparks T C, Thomas J, Watson G B, Zhu Y. 2011. Biological characterization of sulfoxaflor, a novel insecticide. Pest Management Science 67(3): 328-334.
- Bhanu K, Vasanta N, Mallikharjuna M. 2015. Bharata Sulfoxaflor: a new insecticide molecule effective against plant hoppers in rice. Indian Journal of Plant Protection 42(4): 338-342.
- Chandi R S. 2019. Evaluation of new insecticide sulfoxaflor against wheat foliar aphid complex. Journal of Experimental Zoology 22(1): 583-587.
- Deol G S, Gill K S, Brar J S. 1987. Aphid outbreak on wheat and barley in Punjab. Newsletter Aphidological Society of India 6: 7-9.
- Gomez K A, Gomez A A. 1984. Statistical procedures for agricultural research. John Wiley and Sons, USA. 680 pp.
- Jasrotia P, Katare S, Kashyap P L, Kumar S, Kharub A S, Singh G P. 2018. Efficacy of few selected insecticides for the management of corn leaf aphid in barley. Wheat and Barley Research 10(3): 228-30.
- Jasrotia P, Bhardwaj A K, Katare S, Yadav J, Kashyap P L, Kumar S, Singh G P. 2021. Tillage intensity influences insect-pest and predator dynamics of wheat crop grown under different conservation agriculture practices in rice-wheat cropping system of Indo-Gangetic plain. Agronomy 11(6): 1087.
- Katare S, Patel A A, Archarya S. 2015. Bio-efficacy of some new insecticides against foliage feeding barley aphids (*Rhopalosiphum maidis*). Journal of Wheat Research 7(2): 29-33.
- Katare S, Singh B, Patil S. D, Tiwari R, Jasrotia P, Saharan M S, Sharma I. 2017. Evaluation of new insecticides against wheat foliar aphid complex. Indian Journal of Entomology 79(2): 185-190.
- Katare S, Reza, W, Jasrotia P, Saharan M S, Sharma, I. 2018. Efficacy of botanicals and biopesticides against foliage feeding aphids in wheat. Indian Journal of Entomology 80(1): 40-46.
- Kumar A, Singh R, Verma R P S, Singh S. 2001. Barley production technology, Extension Bulletin-11. Directorate of Wheat Research, Karnal.
- Longhurst C, Babcock J M, Denholm I, Gorman K, Thomas J D, Sparks T C. 2013. Cross-resistance relationships of the sulfoximine insecticide sulfoxaflor with neonicotinoids and other insecticides in the whiteflies *Bemisia tabaci* and *Trialeurodes vaporariorum*. Pest Management Science 69(7): 809-813.
- Ozder N. 2002. Development and fecundity of *Sitobion avenae* (F.) (Hom.: Aphididae) on some wheat cultivars in laboratory conditions. Plant Pathology Journal 1(1): 9-10.
- Singh C. 1983. Modern techniques of raising field crops. Oxford and IBH publishing, New Delhi, India.
- Singh B, Deol G S. 2003. Quantitative grain yield losses caused by aphid complex in wheat. Crop Research Hisar 26(3): 501-504.
- Singh B, Jasrotia P. 2020. Impact of integrated pest management (IPM) module on major insect-pests of wheat and their natural enemies in North-western plains of India. Journal of Cereal Research 12(2): 114-119.
- Trdan S, Milevoj L. 1999. The cereal aphid (*Sitobion avenae* F.) wheat pest. Sodobno kmetijstvo (Slovenia) 32(3): 119-128.
- Xin J J, Yu W X, Yi X Q, Gao J P, Gao X W, Zeng X P. (2019). Sublethal effects of sulfoxaflor on the fitness of two species of wheat aphids, *Sitobion avenae* (F.) and *Rhopalosiphum padi* (L.). Journal of Integrative Agriculture 18(7): 1613-1623.
- Yadav R. 2003. A combined source of resistance against corn leaf aphid and yellow rust in barley. International Journal of Pest Management 49(4): 293-296.
- Yu X, Zhang S, Song S, Zhu Q, Xie C, Ji J, Gao X. 2016. Resistance of wheat aphids to six insecticides and assessment of their field efficacy. Acta Entomologica Sinica 59(11): 1206-1212.

(Manuscript Received: May, 2021; Revised: July, 2021;

Accepted: July, 2021; Online Published: August, 2021)

Online published (Preview) in www.entosocindia.org Ref. No. e21090