



## EVALUATION OF INSECTICIDES AND BIOPESTICIDES AGAINST LEAF HOPPER *EMPOASCA KERRI* PRUTHI IN PIGEON PEA

S S KARABHANTANAL\* AND SAICHARAN DHARAVATH<sup>1</sup>

All India Coordinated Sorghum Improvement Project,  
Regional Agricultural Research Station, Vijayapur 586101, Karnataka, India

<sup>1</sup>Department of Agricultural Entomology,  
UAS Dharwad, College of Agriculture, Vijayapur 586101, Karnataka, India

\*Email: sskarabu@gmail.com (ccorresponding author)

### ABSTRACT

Green leafhopper *Empoasca kerri* (Hemiptera: Cicadellidae) is a polyphagous xylem feeder and it is a pest of pulse crops. The present study evaluated some insecticides and biopesticides against *E. kerri* in pigeonpea. The results revealed that its incidence was significantly reduced with flonicamid 50WG and tolfenpyrad 20%SC (89.39 and 81.92%, respectively) after third spray. Also, increase in yield was observed with flonicamid 50WG (13.20 q/ ha) and thiamethoxam 25WDG (12.90 q/ ha) amounting to 37.50 and 36.05%, respectively. The biopesticides viz., neemazol 1% @ 2ml/l and *Lecanicillium lecanii* @2g/l were found to be the least effective.

**Key words:** *Empoasca kerri*, pigeonpea, flonicamid 50WG, tolfenpyrad 20%SC, *Lecanicillium lecanii*, neemazol, thiamethoxam, yield, incidence

In India pigeon pea (*Cajanus cajan* L.) is cultivated in 4.78 million ha with a production of 0.98 mt (Anon, 2019). The insect pests are the main factors for the low productivity and about 250 insect pests under 8 orders and 61 families infest the crop. Since 2012, both nymphs and adults of green leafhopper *Empoasca kerri* Pruthi has been observed to be a major sucking pest and spreading to larger area on pigeonpea. The damage causes phytotoxicity, with hopper burn and economic loss (Singh et al., 2008). This study evaluates insecticides and biopesticides against this pest.

### MATERIALS AND METHODS

Field experiment was conducted at the All India Coordinated Sorghum Improvement Project (AICSP), Regional Agricultural Research Station (RARS), Vijayapur, Karnataka during kharif, 2019. Insecticides including botanicals and entomopathogenic fungi were evaluated in randomized block design (RBD) with eleven treatments replicated thrice. The treatments include- flonicamid 50WG, tolfenpyrad 20%SC, fipronil 5EC, thiamethoxam 25WDG, acephate 75SP, dimethoate 30EC, monocrotophos 36SL, neemazol 1%, buprofezin 20SC, *Lecanicillium lecanii* along with untreated control. Pigeonpea variety TS3R was sown in 33 plots of 5.4x 3.6 m size with a spacing of 90x 30 cm between rows and plants. The crop was raised under rainfed conditions with only one protective irrigation

during the flowering stage. All the recommended agronomic practices were followed. Three sprays were adopted at vegetative and flowering stages, and counts of leafhoppers from three top leaflets were made four times for each spray, as precount, and at 5, 10, and 15 DAT (days after treatment). The data were subjected to corrections after Abbott (1925) before statistical analysis.

### RESULTS AND DISCUSSION

The results revealed that all the treatments exhibited superiority over control in terms of suppressing the leafhopper incidence; flonicamid 50WG @ 100g.a.i./ha resulted in 89.37% reduction over control followed by tolfenpyrad 20%SC @ 200g.a.i./ha with 81.92% reduction. After three sprays the treatments again, flonicamid and tolfenpyrad were the best followed by thiamethoxam 25WDG. The biopesticides *Lecanicillium lecanii* and neemazol 1% resulted in only 46.73 and 41.13% reduction. These results are similar to those of Sunil et al. (2019) that biopesticides *Metarhizium anisopliae* 1.15 WP and NSKE (Neem Seed Kernel Extract) were the least effective. Flonicamid 50WG was found to result in 93.5% reduction in leafhoppers (Duraimurugan and Alivelu, 2017). Anandmurthy et al. (2017) showed that flonicamid 50 WG @ 0.02% gave efficient leafhopper mortality with two sprays. Ram et al. (2020) also reported that thiamethoxam 25% WG

Table 1. Efficacy of insecticides and biopesticides against *E. kerri* in pigeonpea

Treatment	Dosage	Leaf hoppers / 3 top leaflets Pretreatment	15 Days after first application	15 Days after second application	15 Days after third application	% reduction over control after third application	Visual symptom (1-5 Scale)	Yield (q/ ha)
<i>Lecanicillium lecanii</i>	2 g/ l	25.27 (5.08)	19.80 (4.51)	10.27 (3.28)	5.47 (2.44)	46.73	4.67	9.10
Bufrofezin 20SC	200 g.a.i./ ha	23.60 (4.91)	10.87 (3.37)	4.47 (2.23)	1.87 (1.54)	58.16	2.33	10.30
Neemazol 1%	2 ml/l	28.43 (5.38)	18.40 (4.35)	14.27 (3.84)	8.40 (2.98)	41.13	4.67	9.90
Flonicamid 50WG	100 g.a.i./ ha	27.51 (5.29)	8.20 (2.95)	1.60 (1.45)	0.17 (0.82)	89.37	1.33	13.20
Tolfenpyrad 20%SC	200 g.a.i./ ha	24.60 (5.01)	10.03 (3.24)	2.60 (1.76)	0.47 (0.98)	81.92	1.67	12.40
Fipronil 5EC	50 g.a.i./ ha	25.80 (5.13)	12.20 (3.56)	5.27 (2.40)	1.93 (1.56)	63.37	2.33	10.40
Thiamethoxam 25WDG	50 g.a.i./ ha	21.00 (4.64)	9.67 (3.19)	1.80 (1.52)	0.33 (0.91)	81.66	1.33	12.90
Accephate 75SP	750 g.a.i./ ha	24.20 (4.97)	15.27 (3.97)	7.20 (2.77)	4.00 (2.12)	44.44	2.67	10.90
Dimethoate 30EC	600 g.a.i./ ha	27.33 (5.28)	13.67 (3.76)	5.40 (2.43)	1.80 (1.52)	66.66	2.00	11.30
Monocrotophos 36SL	360 g.a.i./ ha	24.47 (5.00)	14.40 (3.86)	7.80 (2.88)	3.80 (2.07)	51.28	2.67	10.60
Untreated control		24.00 (4.95)	26.40 (5.19)	28.03 (5.34)	29.60 (5.49)		5.00	8.25
	C.D (p=0.05)	NS	0.32	0.39	0.24			0.51
	S. Em±	0.28	0.11	0.13	0.08			0.17
	CV (%)	15.20	18.41	13.45	14.66			16.33

\*- Values in parentheses are sine values

(0.5 g/l) reduced leafhoppers after 7 days of spray. The results on the yield indicated that maximum pod yield was obtained with flonicamid 50WG and thiamethoxam 25WDG (13.20 and 12.90 q/ ha, respectively). These results on yield are in accordance with those of Anandmurthy et al. (2017). Chaudhari et al. (2015) also reported increased yield with flonicamid 50WG in Indian bean. Sunil et al. (2019) in groundnut found that thiamethoxam 25WG has shown good yield. Ram et al. (2020) with thiamethoxam 25% WG observed more yields (Table 1).

#### REFERENCES

- Abbott W S. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265-267.
- Anonymous. 2019. *Agricultural Statistics at a Glance*. Directorate of Economics and Statistics, Ministry of Agriculture. Directorate of Economics and Statistics. <http://eands.dacnet.nic.in>
- Anandmurthy T, Parmar G M, Arvindarajan G. 2017. Bio-efficacy of new molecules against sucking pests in summer cowpea. *International Journal of Plant Protection* 10(2): 236-240.
- Chaudhari A J, Korat D M, Dabhi M R. 2015. Bioefficacy of newer insecticides against major insect pests of Indian bean, *Lablab purpureus* L. *Karnataka Journal of Agricultural Sciences* 28: 616-619.
- Duraimurugan P, Alivelu K. 2017. Field efficacy of newer insecticides against sucking insect pests in castor. *Indian Journal of Plant Protection* 45(3): 246-250.
- Ram K M, Ravindra K S, Uadal M, Manohari Lal. 2020. Bio-efficacy of different insecticides against leaf hopper, *Empoasca kerri* Pruthi (Cicadellidae: Hemiptera) in cluster bean. *International Journal of Plant Protection* 13(1): 24-29.
- Sunil G, Sarfraz A, Rohit K N. 2019. Comparative bio-efficacy of newer insecticides and bio-pesticides against leafhopper, *Empoasca kerri* Pruthi of groundnut in semi-arid region of India. *Journal of Entomology and Zoology Studies* 7(5): 112-115.

(Manuscript Received: March, 2021; Revised: August, 2021;  
Accepted: September, 2021; Online Published: November, 2021)  
Online published (Preview) in [www.entosocindia.org](http://www.entosocindia.org) Ref. No. e21093