



## EVALUATION OF INSECTICIDES AGAINST SUCKING PESTS OF INDIAN BEAN

DEVASHRAYEE VAIDIK M, D R PATEL<sup>1</sup> AND P M SANKHLA<sup>1</sup>

Department of Entomology, N M College of Agriculture,  
Navsari Agricultural University (NAU), Navsari 396450, Gujarat, India

<sup>1</sup>Department of Entomology, College of Agriculture, NAU, Bharuch 392012, Gujarat, India  
Email: patel.devendra2829@yahoo.com (corresponding author)

### ABSTRACT

Among the various insecticides evaluated for their field efficacy against sucking pests of Indian bean, the treatments of thiamethoxam 25WG at 0.025%, acetamiprid 20SP at 0.004% and buprofezin 25SC at 0.05% were found to be the most effective against *Aphis craccivora* Koch, *Empoasca kerri* Pruthi and *Bemisia tabaci* Gennadius. While, thiacloprid 21.7SC at 0.012% emerged as the moderately effective. Emamectin benzoate 5SG at 0.002%, indoxacarb 14.5SC at 0.007%, lambda-cyhalothrin 5SC at 0.005% and novaluron 10EC at 0.01% were found to be less effective.

**Key words:** *Lablab purpureus*, *Aphis craccivora*, *Empoasca kerri*, *Bemisia tabaci*, acetamiprid, thiamethoxam, buprofezin, thiacloprid, emamectin benzoate, indoxacarb, lambda-cyhalothrin, novaluron

Indian bean *Lablab purpureus* L. is a legume crop widely grown as vegetable or pulse. Insect pests are major constraints in its productivity. It is attacked by a number of insect pests viz., aphid *Aphis craccivora* Koch.; jassids *Empoasca fabae* (Harris); *E. krameri* Ross and Moore and *E. kerri* Pruthi; pod borer *Etiaella zinckenella* (Treit.); whitefly *Bemisia tabaci* (Genn.); stem fly, *Ophiomyia phaseoli* (Tryon); hairy caterpillars *Ascotis imparta* (Walk.); Bihar hairy caterpillar, *Spilosoma obliqua* (Walk.) etc. Among these, aphids, jassids and whiteflies are the major sucking pests. These pests attack all parts of the plants including pods which result in stunted growth and decreased yield. The honey dew secretion of the aphids provides a suitable media for the development of sooty mould and fungi which ultimately hamper the process of photosynthesis (David and Kumarswami, 1982). Chemical control of *A. craccivora*, *E. kerri* and *B. tabaci* is usually recommended (Garhwal et al., 1994; Dhamaniya et al., 2005; Yadav et al., 2011) but, due to its continuous and enormous use, problems of resistance, deleterious effect on parasitoids and predators, and residue hazards, and environment pollution have arisen. Hence, there is renewed interest in search for new insecticides, and the present study evaluates some of these.

### MATERIALS AND METHODS

The field experiments on the evaluation of efficacy of insecticides were conducted at the College Farm, N M College of Agriculture, Navsari Agricultural University, Navsari, Gujarat during 2019-20. The

variety GNIB-22 was used and the seeds sown in plots of size 11 m<sup>2</sup> at 60x 30 cm spacing in 2<sup>nd</sup> fortnight of October. Eight insecticides viz., thiamethoxam 25WG (1 g/l), thiacloprid 21.7SC (0.6 ml/l), buprofezin 25SC (2.0 ml/l), acetamiprid 20SP (0.2 g/l), indoxacarb 14.5SC (0.5 ml/l), emamectin benzoate 5SG (0.4g/l), lambda-cyhalothrin 5SC (1ml/l) and novaluron 10EC (1 ml/l) were evaluated along with untreated control with each replicated thrice. These were applied as a foliar spray on the crop using pre-calibrated knapsack sprayer when the pest incidence was sufficiently built up. Second spray was repeated after 15 days of the first spray. The observations were recorded a day before spray as well as 1<sup>st</sup>, 3<sup>rd</sup>, 5<sup>th</sup>, 7<sup>th</sup> and 14<sup>th</sup> days after each spray. The observations were made from five randomly selected plants/ plot. *Aphis craccivora* incidence was observed on three randomly selected twigs (about 10 cm in length). *Empoasca kerri* and *B. tabaci* were counted from three leaves (from top, middle and bottom). The pooled data were subjected to statistical analysis.

### RESULTS AND DISCUSSION

The pooled results over two sprays given in Tables 1-3 reveal that significantly minimum incidence of *A. craccivora* was observed in plots treated with thiamethoxam 25WG (3.57 aphids/ twig) and it was at par with acetamiprid 20SP (3.82 aphids/ twig) and buprofezin 25SC (4.04 aphids/ twig), while novaluron 10EC (12.49 aphids/ twig) was found less effective (Table 1). Chaudhary et al. (2015) reported that

Table 1. Efficacy of insecticides against *Aphis craccivora* on Indian bean

Tr. No.	Treatments	Before spray	Mean no. of aphids/ twig														Pooled
			First spray							Second spray							
			IDAS	3DAS	5DAS	7DAS	14DAS	IDAS	3DAS	5DAS	7DAS	14DAS	IDAS	3DAS	5DAS	7DAS	
T <sub>1</sub>	Thiamethoxam 25WG at 0.025%	4.05 (16.43)	2.33 (5.44)	2.23 (4.99)	2.05 (4.20)	1.85 (3.42)	2.26 (5.09)	1.94 (3.75)	1.88 (3.52)	1.66 (2.75)	1.39 (1.95)	1.08 (1.16)	1.89 (3.57)				
T <sub>2</sub>	Thiacloprid 21.7SC at 0.012%	4.13 (17.03)	3.01 (9.04)	2.96 (8.74)	2.84 (8.08)	2.65 (7.00)	2.94 (8.66)	2.68 (7.16)	2.54 (6.48)	2.36 (5.57)	2.16 (4.69)	1.84 (3.38)	2.61 (6.84)				
T <sub>3</sub>	Buprofezin 25SC at 0.05%	3.97 (15.79)	2.41 (5.82)	2.35 (5.54)	2.20 (4.86)	2.02 (4.08)	2.35 (5.52)	2.02 (4.08)	1.91 (3.63)	1.78 (3.19)	1.58 (2.48)	1.25 (1.55)	2.01 (4.04)				
T <sub>4</sub>	Acetamiprid 20SP at 0.004%	4.12 (16.95)	2.37 (5.62)	2.28 (5.19)	2.13 (4.56)	1.94 (3.79)	2.30 (5.27)	1.98 (3.92)	1.89 (3.57)	1.77 (3.12)	1.54 (2.36)	1.13 (1.27)	1.95 (3.82)				
T <sub>5</sub>	Indoxacarb 14.5SC at 0.007%	3.95 (15.58)	3.73 (13.89)	3.68 (13.57)	3.59 (12.91)	3.53 (12.46)	3.58 (12.79)	3.36 (11.31)	3.29 (10.84)	3.13 (9.83)	3.00 (8.98)	2.70 (7.30)	3.37 (11.39)				
T <sub>6</sub>	Emamectin Benzoate 5SG at 0.002%	3.84 (14.72)	3.63 (13.18)	3.59 (12.88)	3.45 (11.94)	3.35 (11.22)	3.52 (12.36)	3.30 (10.87)	3.22 (10.36)	3.08 (9.46)	2.92 (8.50)	2.42 (5.85)	3.26 (10.63)				
T <sub>7</sub>	Lambda-cyhalothrin 5SC at 0.005%	3.97 (15.76)	3.67 (13.51)	3.63 (13.18)	3.53 (12.46)	3.38 (11.44)	3.54 (12.56)	3.33 (11.08)	3.24 (10.54)	3.10 (9.61)	2.94 (8.66)	2.66 (7.07)	3.31 (11.01)				
T <sub>8</sub>	Novaluron 10EC at 0.01%	4.05 (16.40)	4.07 (16.56)	3.92 (15.41)	3.82 (14.61)	3.70 (13.73)	3.63 (13.20)	3.43 (11.81)	3.44 (11.85)	3.21 (10.34)	3.09 (9.56)	2.87 (8.23)	3.53 (12.49)				
T <sub>9</sub>	Control (Treated with water)	4.12 (16.97)	4.65 (21.63)	4.55 (20.67)	4.38 (19.21)	4.22 (17.86)	4.21 (17.76)	3.99 (15.94)	3.94 (15.55)	3.79 (14.38)	3.56 (12.72)	3.27 (10.69)	4.07 (16.56)				
	S.E.m ±	0.19	0.18	0.19	0.16	0.14	0.18	0.17	0.14	0.17	0.14	0.13	0.04				
	S.E.m ± (P×T)	-	-	-	-	-	-	-	-	-	-	-	0.06				
	C.D (p= 0.05)	NS	0.55	0.57	0.48	0.44	0.55	0.51	0.41	0.52	0.43	0.38	0.13				
	C.D (p= 0.05) (P×T)	-	-	-	-	-	-	-	-	-	-	-	NS				

DAS = Days after spraying; Figure in parentheses original, and outside  $\sqrt{x} + 0.5$  transformed values.

Table 2. Efficacy of insecticides against *Empoasca kerri* on Indian bean

Tr. No.	Treatments	Before spray	Mean no. of jassids/ leaf												Pooled
			First spray			Second spray			Pooled						
			1DAS	3DAS	5DAS	7DAS	14DAS	1DAS	3DAS	5DAS	7DAS	14DAS			
T <sub>1</sub>	Thiamethoxam 25WG at 0.025%	2.93 (8.58)	1.91 (3.65)	1.95 (3.82)	1.88 (3.55)	1.73 (2.98)	1.73 (2.98)	1.92 (3.67)	1.68 (2.83)	1.73 (2.99)	1.56 (2.42)	1.17 (1.36)	0.95 (0.90)	1.67 (2.80)	
T <sub>2</sub>	Thiacloprid 21.7SC at 0.012%	2.91 (8.47)	2.32 (5.37)	2.37 (5.63)	2.31 (5.35)	2.22 (4.94)	2.22 (4.94)	2.31 (5.35)	2.17 (4.71)	2.19 (4.82)	2.04 (4.15)	1.79 (3.22)	1.49 (2.22)	2.14 (4.60)	
T <sub>3</sub>	Buprofezin 25SC at 0.05%	2.86 (8.16)	2.30 (5.29)	2.34 (5.47)	2.29 (5.28)	2.20 (4.86)	2.20 (4.86)	2.29 (5.25)	2.13 (4.55)	2.17 (4.72)	1.95 (3.82)	1.72 (2.96)	1.38 (1.90)	2.10 (4.41)	
T <sub>4</sub>	Acetamiprid 20SP at 0.004%	2.83 (7.99)	1.89 (3.56)	1.91 (3.67)	1.84 (3.39)	1.71 (2.92)	1.71 (2.92)	1.88 (3.56)	1.66 (2.76)	1.69 (2.85)	1.51 (2.29)	1.10 (1.22)	0.88 (0.77)	1.63 (2.67)	
T <sub>5</sub>	Indoxacarb 14.5SC at 0.007%	2.88 (8.29)	2.74 (7.51)	2.78 (7.74)	2.74 (7.54)	2.70 (7.27)	2.70 (7.27)	2.74 (7.52)	2.63 (6.92)	2.67 (7.12)	2.52 (6.35)	2.31 (5.34)	2.09 (4.38)	2.61 (6.82)	
T <sub>6</sub>	Emamectin Benzoate 5SG at 0.002%	2.90 (8.39)	2.71 (7.36)	2.74 (7.54)	2.71 (7.34)	2.66 (7.06)	2.66 (7.06)	2.72 (7.41)	2.60 (6.78)	2.65 (7.02)	2.47 (6.08)	2.30 (5.29)	2.09 (4.35)	2.58 (6.67)	
T <sub>7</sub>	Lambda-cyhalothrin 5SC at 0.005%	2.82 (7.95)	2.77 (7.65)	2.80 (7.87)	2.75 (7.59)	2.71 (7.34)	2.71 (7.34)	2.78 (7.73)	2.65 (7.04)	2.71 (7.34)	2.56 (6.55)	2.36 (5.59)	2.15 (4.64)	2.64 (6.99)	
T <sub>8</sub>	Novaluron 10EC at 0.01%	2.87 (8.23)	2.80 (7.86)	2.83 (8.00)	2.78 (7.75)	2.75 (7.56)	2.75 (7.56)	2.80 (7.88)	2.69 (7.23)	2.73 (7.49)	2.56 (6.55)	2.41 (5.81)	2.19 (4.80)	2.67 (7.15)	
T <sub>9</sub>	Control (Treated with water)	2.84 (8.10)	2.99 (8.96)	3.03 (9.16)	2.99 (8.92)	2.94 (8.64)	2.94 (8.64)	2.99 (8.98)	2.84 (8.08)	2.88 (8.34)	2.86 (8.20)	2.69 (7.25)	2.53 (6.40)	2.89 (8.36)	
	S.E.m ±	0.12	0.13	0.11	0.12	0.14	0.14	0.12	0.14	0.12	0.09	0.11	0.10	0.03	
	S.E.m ± (P×T)	-	-	-	-	-	-	-	-	-	-	-	-	0.05	
	C.D (p= 0.05)	NS	0.39	0.34	0.36	0.40	0.40	0.37	0.41	0.36	0.27	0.34	0.32	0.09	
	C.D (p= 0.05) (P×T)	-	-	-	-	-	-	-	-	-	-	-	-	NS	

DAS = Days after spraying; Figure in parentheses are original value whereas, those outside are  $\sqrt{x + 0.5}$  transformed values.

Table 3. Efficacy of insecticides against *Bemisia tabaci* on Indian bean

Tr. No.	Treatments	Before spray	Mean no. of whitefly/ leaf												Pooled			
			First spray				Second spray				Pooled							
			1DAS	3DAS	5DAS	7DAS	14DAS	IDAS	3DAS	5DAS	7DAS	14DAS	IDAS	3DAS	5DAS	7DAS	14DAS	
T <sub>1</sub>	Thiamethoxam 25WG at 0.025%	2.73 (7.45)	2.11 (4.47)	2.12 (4.48)	1.84 (3.39)	1.63 (2.68)	1.86 (3.46)	1.47 (2.16)	1.48 (2.18)	1.39 (1.92)	1.18 (1.39)	1.01 (1.03)	1.47 (2.16)	1.48 (2.18)	1.39 (1.92)	1.18 (1.39)	1.01 (1.03)	1.63 (2.68)
T <sub>2</sub>	Thiacloprid 21.7SC at 0.012%	2.68 (7.18)	2.53 (6.40)	2.56 (6.55)	2.31 (5.35)	2.13 (4.54)	2.30 (5.27)	1.99 (3.97)	2.00 (4.01)	1.92 (3.67)	1.74 (3.03)	1.54 (2.36)	1.99 (3.97)	2.00 (4.01)	1.92 (3.67)	1.74 (3.03)	1.54 (2.36)	2.12 (4.51)
T <sub>3</sub>	Buprofezin 25SC at 0.05%	2.75 (7.54)	2.14 (4.59)	2.14 (4.58)	1.86 (3.47)	1.70 (2.89)	1.90 (3.62)	1.50 (2.24)	1.52 (2.30)	1.45 (2.10)	1.22 (1.48)	1.07 (1.15)	1.50 (2.24)	1.52 (2.30)	1.45 (2.10)	1.22 (1.48)	1.07 (1.15)	1.67 (2.82)
T <sub>4</sub>	Acetamiprid 20SP at 0.004%	2.71 (7.36)	2.09 (4.38)	2.10 (4.41)	1.83 (3.36)	1.62 (2.62)	1.83 (3.36)	1.40 (1.96)	1.43 (2.04)	1.33 (1.77)	1.16 (1.34)	0.94 (0.90)	1.40 (1.96)	1.43 (2.04)	1.33 (1.77)	1.16 (1.34)	0.94 (0.90)	1.60 (2.57)
T <sub>5</sub>	Indoxacarb 14.5SC at 0.007%	2.82 (7.97)	2.73 (7.43)	2.79 (7.80)	2.65 (7.05)	2.54 (6.47)	2.63 (6.93)	2.40 (5.76)	2.43 (5.92)	2.32 (5.37)	2.11 (4.47)	1.92 (3.67)	2.40 (5.76)	2.43 (5.92)	2.32 (5.37)	2.11 (4.47)	1.92 (3.67)	2.47 (6.11)
T <sub>6</sub>	Emamectin Benzoate 5SG at 0.002%	2.76 (7.64)	2.72 (7.38)	2.77 (7.69)	2.63 (6.93)	2.53 (6.40)	2.62 (6.88)	2.38 (5.66)	2.42 (5.84)	2.32 (5.37)	2.10 (4.42)	1.89 (3.56)	2.38 (5.66)	2.42 (5.84)	2.32 (5.37)	2.10 (4.42)	1.89 (3.56)	2.45 (6.03)
T <sub>7</sub>	Lambda-cyhalothrin 5SC at 0.005%	2.80 (7.84)	2.76 (7.60)	2.81 (7.93)	2.67 (7.12)	2.58 (6.67)	2.64 (7.00)	2.45 (6.03)	2.51 (6.28)	2.35 (5.54)	2.16 (4.65)	2.00 (4.00)	2.45 (6.03)	2.51 (6.28)	2.35 (5.54)	2.16 (4.65)	2.00 (4.00)	2.51 (6.31)
T <sub>8</sub>	Novaluron 10EC at 0.01%	2.70 (7.30)	2.76 (7.65)	2.83 (8.00)	2.72 (7.41)	2.61 (6.83)	2.68 (7.16)	2.47 (6.11)	2.52 (6.37)	2.37 (5.62)	2.21 (4.87)	2.05 (4.22)	2.47 (6.11)	2.52 (6.37)	2.37 (5.62)	2.21 (4.87)	2.05 (4.22)	2.54 (6.46)
T <sub>9</sub>	Control (Treated with water)	2.79 (7.78)	2.90 (8.45)	3.01 (9.08)	3.04 (9.24)	2.96 (8.47)	3.00 (9.00)	2.85 (8.16)	2.89 (8.39)	2.82 (7.95)	2.71 (7.34)	2.46 (6.04)	2.85 (8.16)	2.89 (8.39)	2.82 (7.95)	2.71 (7.34)	2.46 (6.04)	2.88 (8.31)
	S.E.m ±	0.13	0.13	0.12	0.10	0.10	0.10	0.12	0.12	0.13	0.10	0.08	0.12	0.12	0.13	0.10	0.08	0.06
	S.E.m ± (P×T)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.04
	C.D (p= 0.05)	NS	0.38	0.36	0.32	0.32	0.30	0.37	0.37	0.38	0.31	0.25	0.37	0.37	0.38	0.31	0.25	0.20
	C.D at 5 % (P×T)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.12

DAS = Days after spraying; Figure in parentheses are original value whereas, those outside are  $\sqrt{x + 0.5}$  transformed values.

imidacloprid followed by acetamiprid were superior; Choudhary et al. (2017) showed that thiamethoxam (0.005%) was the most effective. As regards the incidence of *E. kerri*, acetamiprid 20SP (2.67 jassids/ leaf) followed by thiamethoxam 25WG (2.80 jassids/ leaf) were the best (Table 2); buprofezin 25SC and thiacloprid 21.7SC were the next best. Chaudhary et al. (2015) found that imidacloprid followed by acetamiprid were superior; while Singh et al. (2019) showed that acetamiprid (0.004%) was the most effective in green gram. Meena et al. (2020) reported that imidacloprid > thiamethoxam > acetamiprid were the most effective in green gram. The incidence of *B. tabaci* was significantly reduced with acetamiprid 20SP (2.57 whiteflies/ leaf) and it was at par with thiamethoxam 25WG (2.68 whiteflies/ leaf); it was followed by buprofezin 25SC and thiacloprid, while was the least effective (Table 3). These results agree with those of Chaudhary et al. (2015) that incidence of whitefly was significantly reduced with acetamiprid. Kukvaya et al. (2018) revealed that thiamethoxam 25 WG @ 0.005 was highly effective against whitefly in moth bean. Singh and Singh (2018, 2019) observed that acetamiprid was the most effective in green gram.

#### REFERENCES

Chaudhari A J, Korat D M, Dabhi M R. 2015. Bioefficacy of newer insecticides against major insect pests of Indian bean, *Lablab*

*purpureus* L. Karnataka Journal of Agricultural Sciences 28(4): 616-619.

Choudhary A L, Hussain A, Choudhary M D, Samota R, Jat S. 2017. Bio-efficacy of newer insecticides against aphid, *Aphis craccivora* Koch on cowpea. Journal of Pharmacognosy and Phytochemistry 6(4): 1788-1792.

David B V, Kumaraswami T. 1982. Elements of Economic Entomology Popular Book Depot, Madras, 173 pp.

Dhamaniya B, Sharma J K, Kumawat K C. 2005. Bio-efficacy of insecticides against sucking insect pests of moth bean, *Vigna aconitifolia*. Annals of Plant Protection Sciences 13(1): 91-93.

Garhwal S N, Verma S K, Sharma J K. 1994. Field efficacy of different insecticide against cowpea aphid, *Aphis craccivora* Koch. Annals of Arid Zone 33(2): 159-160.

Kukvaya D D, Jakhar B L, Chaudhari S J, Patel B C. 2018. Bio-efficacy of insecticides against sucking insect pests of moth bean, *Vigna aconitifolia* (Jacq.) Marechal. Journal of Entomology and Zoology Studies 6(5): 2227-2230.

Meena R K, Meena R K, Singh Uadal, Meena M L, Meena B L, Anokhe A. 2020. Evaluation of insecticides against leaf hopper *Empoasca kerri* Pruthi in green gram. Indian journal of Entomology 82(4): 768-770.

Singh M, Bairwa D K, Dadrwal B K, Chauhan J. 2019. Relative efficacy of new generation insecticides against sucking insect pests of green gram. Journal of Pharmacognosy and Phytochemistry 8(2): 882-886.

Singh S K, Singh P S. 2018. Efficacy of some insecticides and biopesticides against whitefly, *Bemisia tabaci* on green gram. Journal of Experimental Zoology India 21(1): 241-246.

Yadav S R, Kumawat K C, Khinchi S K. 2011. Efficacy of new insecticide molecules and bioagents against sucking insect pests of cluster bean, *Cyamopsis tetragonoloba* (Linn.) Taub. Journal of Plant Protection and Environment 8(1): 115-122.

(Manuscript Received: July, 2021; Revised: January, 2022;

Accepted: January, 2022; Online Published: March, 2022)

Online published (Preview) in [www.entosocindia.org](http://www.entosocindia.org) Ref. No. e21164