



## SUITABILITY OF BLACK SOLDIER FLY LARVAE AS HOST FOR ENTOMOPATHOGENIC NEMATODES

MANU C R,\* N ANITHA, GAVAS RAGESH<sup>1</sup> AND POORNIMA YADAV P P<sup>2</sup>

Department of Agricultural Entomology, College of Agriculture, Kerala Agricultural University, Vellayani, Thiruvananthapuram 695522, Kerala, India

<sup>1</sup>Banana Research Station, Kerala Agricultural University, Kannara, Thrissur 680652, Kerala, India

<sup>2</sup>Krishi Vigyan Kendra Kollam, Kerala Agricultural University, Sadanandapuram, Kollam 691531, Kerala, India

\*Email: manudasc2014@gmail.com (corresponding author)

### ABSTRACT

Black soldier fly *Hermetia illucens* (L.) larvae can serve many uses, of which potential exists for its use as a host for mass production of entomopathogenic nematodes. This study evaluates its larval instars for their suitability for rearing of *Heterorhabditis bacteriophora* (Poinar) and *Steinernema carpocapsae* (Weiser). The results revealed that 4<sup>th</sup> instar larvae is economical and effective, and this can be used at farm level. Thus, it can be popularized for mass production of entomopathogenic nematodes. The nematodes/ cadaver was found to be half as that from *Galleria*, and hence two 4<sup>th</sup> instar cadavers can be recommended in place of a single *Galleria* cadaver. The results also showed that *H. illucens* is unsuitable for mass multiplication of *S. carpocapsae*.

**Key words:** *Hermetia illucens*, *Heterorhabditis bacteriophora*, *Steinernema carpocapsae*, *Galleria mellonella*, insect rearing, mass production, larval instars, cadavers

Black soldier fly *Hermetia illucens* (L.) (Stratiomyidae: Diptera) is a beneficial insect as it can be used for waste management with minimum carbon footprint, besides its other uses. It can be easily reared in any organic waste material and its biology had been studied by Sharanabasappa et al. (2019), and with biological parameters compared on different food wastes (Srikanth and Deshmukh, 2021). The present study assessed the use of *H. illucens* larva as host for the rearing of entomopathogenic nematodes. These nematodes (EPN) are being advocated as the safest biopesticides. Entomopathogenic nematodes of genera *Steinernema* and *Heterorhabditis* are efficient biocontrol agents (Koppenhofer, 2000). Nematode infected cadavers of greater wax moth (*Galleria mellonella* L.) larvae are the popular ones for the rearing of these nematodes, but their rearing and maintenance requires sufficient expertise and care. This study evaluates the suitability of *H. illucens* larvae as hosts for cost effective mass multiplication of some entomopathogenic nematodes.

### MATERIALS AND METHODS

The experiment was conducted at the Department of Agricultural Entomology, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala

(N8°25'37.13952", E76°59'12.50736", 29 masl) under laboratory condition during 2020-21. Egg masses of *H. illucens* were collected in cardboard honeycombs from a compost bin set up for attracting the naturally occurring females for oviposition (Ewusie et al., 2019). These were transferred to plastic pots (17 and 25 cm dia, 30cm high) containing vegetable and fruit waste. Suitable larval instars were collected from these and simultaneously reared final instar larvae of *G. mellonella* on standard semisynthetic diet were used as check. Entomopathogenic nematodes viz., *Heterorhabditis bacteriophora* (Poinar) and *Steinernema carpocapsae* (Weiser), obtained from the Banana Research Station, Kannara and ICAR-NBAIR Bengaluru, respectively were used. Nematodes were maintained in the laboratory (28±2°C, 60-70%RH) as aqueous suspension as well as infected cadavers. There were 10 treatments (T1: 5<sup>th</sup> instar BSFL+ *H. bacteriophora*, T2: 4<sup>th</sup> instar BSFL+*H. bacteriophora*, T3: 3<sup>rd</sup> instar BSFL + *H. bacteriophora*, T4: *Galleria* larva+ *H. bacteriophora*, T5: 5<sup>th</sup> instar BSFL+ *S. carpocapsae*, T6: 4<sup>th</sup> instar BSFL+*S. carpocapsae*, T7: 3<sup>rd</sup> instar BSFL + *S. carpocapsae*, T8: *Galleria* larva+ *S. carpocapsae*, T9: 6<sup>th</sup> instar BSFL+ *H. bacteriophora* and T10: 6<sup>th</sup> instar BSFL + *S. carpocapsae*) replicated four times under completely randomized design. One

replication consisted of 10 uniform sized larvae of respective instars in a 90 mm petriplate lined with Whatman number 1 filter paper disc. The larvae were exposed to 100 infective juveniles (IJs) of respective nematodes per larva (Koppenhofer and Kaya 1998).

For this purpose, freshly emerged IJs were collected from Whites trap (White, 1927) in double distilled water and diluted to obtain the required number of IJs; the solution was applied to filter paper disc in petri plate. A waiting period of 10 min was given to ensure uniform distribution of IJs and the uniform soaking of filter paper, larvae were introduced and closed. The petriplates were incubated at laboratory conditions at room temperature. Observations were taken at an interval of 24 hr on mortality. Absolute control was set using double distilled water with above mentioned insects and was also replicated four times. Half of the cadavers formed in various treatments were kept in petri plates in laboratory conditions under room temperature and observed daily for colour change, intactness of cadaver and natural emergence of nematodes. Other half of the cadavers were kept in White trap and number of IJs emerged were observed and counted under stereo zoom microscope on 24 hr basis. Viability of IJs from

BSF larvae were found out by infecting them against *Galleria* larvae. The data obtained were subjected to statistical analysis using the WASP-Web Agri Stat Package 2.0 programme developed by ICAR- Central Coastal Agricultural Research Institute, Goa.

## RESULTS AND DISCUSSION

The results revealed that treatments involving only double distilled water did not cause mortality (these not shown in tables); all larvae were alive in the treatments T5: 5<sup>th</sup> instar BSF + *S. carpocapsae*; T9: 6<sup>th</sup> instar BSF + *H. bacteriophora* and T10: 6<sup>th</sup> instar BSF + *S. carpocapsae*; and larvae in these successfully completed their lifecycle. Mortality of larvae when observed daily for 10 days, it was seen that after 24 hr, mortality was there only in *Galleria* larvae (Table 1). The EPN, *H. bacteriophora* was able to bring about mortality among *H. illucens* larval instars from second day; and all larvae in T4: *Galleria* larva+ *H. bacteriophora* also died on 2 DAI. On the third day, *S. carpocapsae* could bring about 100% mortality of *Galleria* larvae (T8), whereas 100% mortality was observed in T3 (3<sup>rd</sup> instar BSF + *H. bacteriophora*) after 5 days. *Galleria mellonella*, the universal host of entomopathogenic nematodes

Table 1. Mortality of BSF larvae and *Galleria* larvae treated with entomopathogenic nematodes

Treat-ments	Mortality at different days interval (%)										Total
	1	2	3	4	5	6	7	8	9	10	
T1	0 (0.91)c	2.5 (5.29)e	2.5 (5.29)d	5 (9.67)de	5 (9.67)de	5 (9.67)de	5 (9.67)d	5 (9.67)d	5 (9.67)d	5 (9.67)d	5 (9.67)e
T2	0 (0.91)c	37.5 (37.51)d	75 (61.23)b	80 (64.18)b	85 (67.50)b	85 (67.50)b	85 (67.50)b	85 (67.50)b	85 (67.50)b	85 (67.50)b	87.5 (69.53)b
T3	0 (0.91)c	82.5 (65.47)c	82.5 (65.50)b	97.5 (84.71)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a
T4	85 (67.87)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a	100 (89.09)a
T5	0 (0.91)c	0 (0.91)e	0 (0.91)d	7.5 (0.91)e	17.5 (0.91)e	17.5 (0.91)d	40 (0.91)d	45 (0.91)d	45 (0.91)d	45 (0.91)d	35 (0.91)f
T6	0 (0.91)c	2.5 (0.91)e	7.5 (0.91)d	32.5 (11.70)d	42.5 (18.34)d	57.5 (18.34)d	67.5 (38.95)c	75 (42.12)c	80 (42.12)c	82.5 (42.12)c	77.5 (36.22)d
T7	75 (0.91)c	92.5 (5.29)e	100 (14.05)c	100 (34.50)c	100 (40.39)c	100 (50.14)c	100 (59.00)b	100 (63.52)b	100 (66.53)b	100 (68.19)b	100 (61.77)c
T8	0 (60.64)b	0 (78.30)b	0 (89.09)a	0 (89.09)a	0 (89.09)a	0 (89.09)a	0 (89.09)a	0 (89.09)a	0 (89.09)a	0 (89.09)a	0 (89.09)a
T9	0 (0.91)c	0 (0.91)e	0 (0.91)d	0 (0.91)e	0 (0.91)e	0 (0.91)d	0 (0.91)d	0 (0.91)d	0 (0.91)d	0 (0.91)d	0 (0.91)f
T10	0 (0.91)c	0 (0.91)e	0 (0.91)d	0 (0.91)e	0 (0.91)e	0 (0.91)d	0 (0.91)d	0 (0.91)d	0 (0.91)d	0 (0.91)d	0 (0.91)f
CV	27	22.15	17.67	17.91	21.54	22.1	19.16	16.45	14.95	14.37	8.43
CD (0.05)	5.29	9.10	8.37	9.98	12.62	13.27	12.31	10.75	9.84	9.49	5.44
SEM	3.36	9.93	8.39	11.93	19.10	21.10	18.17	13.87	11.61	10.81	1.90

T1: 5<sup>th</sup> instar BSF+ *H. bacteriophora*, T2: 4<sup>th</sup> instar BSF +*H. bacteriophora*, T3: 3<sup>rd</sup> instar BSF + *H. bacteriophora*, T4: *Galleria* larva+ *H. bacteriophora*, T5: 5<sup>th</sup> instar BSF + *S. carpocapsae*, T6: 4<sup>th</sup> instar BSF +*S. carpocapsae*, T7: 3<sup>rd</sup> instar BSF + *S. carpocapsae*, T8: *Galleria* larva+ *S. carpocapsae*, T9: 6<sup>th</sup> instar BSF + *H. bacteriophora* and T10: 6<sup>th</sup> instar BSF + *S. carpocapsae*; \* Values in parentheses after arc sine transformation

proved its superiority. Alonso et al. (2018) found that *S. carpocapsae* activation was much higher to insect homogenates of *G. mellonella* than to *H. illucens*. This is slightly in disagreement with the findings of Tautoris et al. (2017) who observed that fourth instar of *H. illucens* was the most susceptible, and third instar was more prone to speedy nematode infections since both nematodes caused mortality within 48 hr. Later instars of *H. illucens* are known for their tough integument due to deposition of calcium carbonate (Johannsen, 1922); their larvae have only two pairs of functional spiracles and the integument is shagreened due to deposition of calcium carbonate (Stehr, 2008).

Comparing the nematodes, *H. bacteriophora* gave quick mortality than *S. carpocapsae*; the former not only utilized natural openings but it got attached to host cuticle and entered with a drill like mechanism and caused changes in host behavior and produced septicemia within 6 hr in *Drosophila* maggots (Dziedziech et al., 2020). The results clearly indicated that mortality decreased with age of the larvae. A similar trend was also observed by Tourtois et al. (2017). Younger larvae of Mexican fruit fly *Anastrepha ludens* (Loew) were found more susceptible to *H. bacteriophora* (Toledo et al., 2005) than older ones; which is in agreement with the present findings. Susceptibility of younger larvae to EPN was also demonstrated by Fuxa et al. (1988) in the case of *S. feltiae* exposure to fall armyworm larvae.

Number of infective juveniles of EPN that emerged from cadavers were a direct indicator of host suitability for mass multiplication. From the *H. bacteriophora* infected *Galleria* cadaver, 4.51 lakhs of IJs emerged, whereas from 4<sup>th</sup> instar of *H. illucens*, 2.3 lakhs *H. bacteriophora* IJs were seen emerging. Rahoo et al. (2019) also observed >4.5 lakhs *H. bacteriophora* IJs emerging from a single final instar *G. mellonella* cadaver. The present observations in this respect are not in agreement with that of Tourtois et al. (2017) who observed tenfold less IJs of *H. bacteriophora* from *H. illucens* than from *G. mellonella*, even after injuring the larval cuticle. *Hermetia illucens* BSF was observed not at all suitable for *S. carpocapsae* mass production; though 4<sup>th</sup> instar larva produced 1.1 lakh IJs; the emergence was not uniform in all replications. Increased infection rate and mortality did not lead to an increase in the number of IJs harvested from *H. illucens* exposed to *Steinernema* spp. (Tourtois, 2014). In the case of *H. bacteriophora*, IJs started emerging from *G. mellonella* and 4<sup>th</sup> instar of *H. illucens* from the 6<sup>th</sup> day but cease to emerge from 27 and 23<sup>rd</sup> day, respectively




Fig. 1. Daily emergence of infective juveniles from cadavers

(Fig. 1). Peak emergence of *H. bacteriophora* was seen on the 10<sup>th</sup> day on both of these cadavers. A similar trend was observed by Rahoo et al. (2019) who observed a peak on the 13<sup>th</sup> day in the case of *H. bacteriophora* from *Galleria* cadavers. For *S. carpocapsae*, infective juveniles emerged from 6<sup>th</sup> to 16<sup>th</sup> day, 11<sup>th</sup> to 21<sup>st</sup> day and 11<sup>th</sup> to 13<sup>th</sup> day in *Galleria*, third and fourth instar *H. illucens* larvae, respectively; though third instar was more susceptible, these produced less nematodes/insect.

Cadavers formed by *H. bacteriophora* were initially colourless, turning pale yellowish orange within a day, then to brick red and chocolate brown within 4 to 6 days. All cadavers were flaccid initially, but they were seen gaining turgidity slowly within 5 to 7 days. Cadavers formed due to attack of *S. carpocapsae* were initially colourless, flaccid and softer than that of *H. bacteriophora*, turned to pale grey and to dark grey in *Galleria*.; while *H. illucens* cadavers also shown variations in colour from colourless to various shades of grey, but the colour did not get intensified. Natural egress of infective juveniles was observed in the case of cadavers of *G. mellonella* (both nematodes) and *H. bacteriophora* infected 4<sup>th</sup> instar *H. illucens* only. Nematodes were seen emerging from the mouth in the case of *H. illucens* and through all natural openings in *G. mellonella* initially. Infective juveniles of both species of nematodes from *H. illucens* cadavers were found viable since 100% mortality was observed in *Galleria* larvae when inoculated with these IJs emerged from *H. illucens* BSF (Fig. 2). This clearly indicated that EPN emerged from *H. illucens* are as viable and effective as that from *G. mellonella*. Thus, it can be concluded that fourth instar *H. illucens* can be utilized for the mass production of *H. bacteriophora*.




Fig. 2. Mortality of *G. mellonella* larvae by IJs from two hosts

### ACKNOWLEDGEMENTS

Authors acknowledge Kerala Agricultural University for providing the facilities for research.

### FINANCIAL SUPPORT

Authors acknowledge Kerala Agricultural University for the financial support.

### AUTHOR CONTRIBUTION STATEMENT

MCR, NA and GR conceived research. GR helped in designing the research and provided guidance in carrying out nematological techniques and provided nematode culture. PYPI contributed to statistical analysis. MCR prepared the draft manuscript. All authors reviewed, updated and finalized the manuscript. All authors read and approved the manuscript.

### CONFLICTS OF INTEREST / COMPETING INTERESTS

The authors hereby declare that there are no conflicts of interest.

### REFERENCES

- Alonso V, Nasrolahi S, Dillman A R. 2018. MDPI, 10. <https://www.mdpi.com/2075-4450/9/2/59> (20 March 2021).

Dziedzic A, Shivankar S, Theopold U. 2020. MDPI, 15. <https://www.mdpi.com/2075-4450/11/1/60> (20 March 2021).

Ewusie E A, Kwapong P K, Ofosu-Budu G, Sandrock C, Akumah A M, Nartey E K, Tetegaga C, Agyakwah S K. 2019. ScienceDirect, 12. <https://www.sciencedirect.com/science/article/pii/S2468227619306957> (20 March 2021).

Fuxa J R, Richter A R, Acudelo - Silva F. 1988. Effect of host age and nematode strain on susceptibility of *Spodoptera frugiperda* to *Steinernema feltiae*. Journal of Nematology 20: 91-95.

Johannsen O A. 1922. Stratiomyiid larvae and puparia of the North Eastern States. Journal of the New York Entomological Society 30(4): 141-153.

Koppenhofer A M, Kaya H K. 1998. Synergism of imidacloprid and entomopathogenic nematodes: A novel approach to white grub control in turf grass. Journal of Economic Entomology 91(3): 618-623.

Koppenhofer A M. 2000. Nematodes. Lacey L A, Kaya H K (eds.). Field manual of techniques in invertebrate pathology. Kluwer, Academic Press, Dordrecht. 888 pp.

Rahoo A L, Mukhtar T, Bughio B A, Rahoo R K. 2019. Relationship between the size of *Galleria mellonella* larvae and the production of *Steinernema feltiae* and *Heterorhabditis bacteriophora*. Pakistan Journal of Zoology 51(1): 79-84.

Sharanabasappa D, Srikanth B H, Maruthi M S, Pavithra H B. 2019. Biology of black soldier fly *Hermetia illucens* (L.) (Diptera: Stratiomyidae) on musk melon fruit. Indian Journal of Entomology 81(1): 153-155

Srikanth B H, Deshmukh S. 2021. Growth performance and bioconversion rate of black soldier fly *Hermetia illucens* (L.) Indian Journal of Entomology 83(2): 231-234

Stehr F W. 2008. Immature insects- volume 2 (updated printing). Kendall/Hunt Publishing Company, US. 992 pp.

Toledo J, Ibarra J E, Liedo P, Gomez A, Rasgado M A, Williams T. 2005. Infection of *Anastrepha ludens* (Diptera: Tephritidae) larvae by *Heterorhabditis bacteriophora* (Rhabditida: Heterorhabditidae) under laboratory and field conditions. Biocontrol Science and Technology 15(6): 627-634.

Tourtois J, Ali J G, Grieshop M. 2017. Susceptibility of wounded and intact blacksoldier fly *Hermetia illucens* (L.) (Diptera: Stratiomyidae) to entomopathogenic nematodes. Journal of Invertebrate Pathology 150:121-129.

Tourtois J. 2014. On entomopathogenic nematodes (Rhabditida: Steinernematidae and Heterorhabditidae): A potential rearing host, black soldier fly *Hermetia illucens* (L.) (Diptera: Stratiomyidae) and compatibility with a predatory beetle, *Dalotia coriaria* (Kraatz) (Coleoptera: Staphylinidae). M Sc Thesis. Michigan State University. USA. 102 pp.

White G F. 1927. A method for obtaining infective nematode larvae from cultures. Science. 66 (1709): 302-303.

(Manuscript Received: June, 2021; Revised: November, 2021;  
Accepted: December, 2021; Online Published: February, 2022)  
Online published (Preview) in [www.entosocindia.org](http://www.entosocindia.org) Ref. No. e21179