



MYCOPHAGOUS PHLAEOTHRIPIDAE (THYSANOPTERA: TUBULIFERA) IN THE INDIAN SUBCONTINENT

R VARATHARAJAN

Department of Zoology, Centre of Advanced Study in Life Sciences
Manipur University, Imphal 795003, Manipur, India
Email: rvrajanramya@gmail.com

ABSTRACT

Mycophagous Phlaeothripidae (Thysanoptera: Tubulifera:) are recognizable as (a) fungal-hyphae-feeding mycetophagous Phlaeothripinae, living on fungi that infest dry twigs and (b) fungal-spore-feeding sporophagous Idolothripinae, usually present on leaf litter. In the Indian subcontinent, out of 765 species of known Thysanoptera, nearly 152 species in 66 genera fall within the mycophagous group of the Phlaeothripidae with 54 idolothripine and 98 phlaeothripine species. Taxonomic diversity of these species in bamboo- and oak-leaf litter and pine forests in the sub-Himalayan ecosystems of North-Eastern India are discussed in this article, along with the diversity that is apparent in their developmental patterns, thrips-plant-fungus association, and phenotypic plasticity, supplemented with a note on their ecological implications.

Key words: Phlaeothripidae, bamboo, pine, plant fungus, association, development, ecological implications, fungal spores and mycelia, taxonomic diversity, leaf litter, phenotypic plasticity, oak leaf litter

Thrips (Thysanoptera) are relatively small insects with asymmetrical mouthparts bearing fringed wings and protractible, adhesive pre-tarsal bladders. The term *thrips* denotes both singular and plural forms, meaning 'woodworm' in ancient Greek, besides other commonly used terms, such as the thunderflies, storm flies, thunderblights, storm bugs, thunderbugs, corn fleas, corn flies, corn lice, freckle bugs, harvest bugs and physopods (Kirk, 1996; Marren and Mabey, 2010; Kobro, 2011). Most of them feed on plant sap (e.g., *Ananthakrishnana euphorbiae* (Priesner), and some of them on floral nectar and pollen (e.g., *Dichromothrips nakahari* Mound, Thripidae), fungal spores (e.g., *Holurothrips manipurensis* Varatharajan and Chochong) and mycelia (e.g., *Adraneothrips okajimai* (Muraleedharan and Sen)), further to a few species being predatory (e.g., *Androthrips ramachandrai* Karny (all Phlaeothripidae except *D. nakahari*) feeding on other smaller soft-bodied insects and mites. Feeding niches of the Phlaeothripidae vary highly, most feed on plants, and many others feed on pollen, fungal filaments and spores, and some of them feed on other arthropods. Thysanoptera comprises two suborders, the Terebrantia and Tubulifera, of which the body length of those belonging to Terebrantia are always <1 mm, while those of the Tubulifera can be from 1.1 to 15 mm, such as the spore-feeding taxa of the Phlaeothripidae, viz., *Bactrothrips* Karny, *Elaphrothrips*

Buffa, *Mecynothrips* Bagnall, *Meiothrips* Priesner, *Oidanothrips* Moulton, and *Tiarothrips* Priesner (Ananthakrishnan, 1973; Eow et al. 2011; Mound and Tree, 2011) with the exception of the smallest leaf-litter inhabiting *Preeriella* Hood (Phlaeothripinae) being <1 mm (Okajima, 1998). Although considerable volume of information on fungus-feeding Thysanoptera has been documented, the present article refers to the diversity and dynamics of mycophagous Phlaeothripidae of the Indian subcontinent, drawing examples from the North-Eastern Himalayan parts of India.

A. Diversity of the mycophagous Thysanoptera

ThripsWiki (2021) lists 6312 extant taxa under 785 genera and 175 extinct taxa under 65 genera (Mound and Vesmanis, 2021), of which a little >50% utilize fungi as their source of food (Morse and Hoddle, 2006; Mound, 2005). Among the mycophagous Thysanoptera, >700 spore-feeding idolothripines are known worldwide (Eow et al., 2011). In the Neotropics, mycophagous Thysanoptera occupy 50% of the total numbers of thrips known (Mound, 2002). The Uzelothripidae, Merothripidae, and Idolothripinae are primarily fungal-spore feeders, whereas half of the world's known Phlaeothripidae are mycelial feeders. In the Indian subcontinent out of 765 known taxa (Tyagi and Kumar, 2016; Rachana and Varatharajan, 2017), 152 are mycophagous including 54 species of spore-

feeding idolothripines and 98 species of mycetophagous phlaeothripines (Table 1, 2). On extrapolating this statistic, we can assume that close to 1400 species of mycetophagous Thysanoptera occur in the world, since *c.* 700 Idolothripinae are known (Mound et al., 2013). Although the Paraneoptera (Thysanoptera) has been categorized into 15 families including that of the fossil-spore feeding Uzelothripidae (Mound, 2013), members of Phlaeothripidae include *c.* 3500 species (ThripsWiki, 2021), of which approximately 2100 belong to fungus-feeding *Phlaeothrips* lineage (Dang et al., 2014).

In oak-leaf litter

The Sub-Himalayan region stretching from Jammu and Kashmir to Manipur (24°-34°N, 93°-95°E) is the *Quercus* (Fagaceae)-belt of the Indian subcontinent (Negi and Naithani, 1995). *Quercus*-leaf litter harbours at least 14 species of mycophagous Phlaeothripidae, of which five are idolothripines and the remainder are phlaeothripines, which dominate both in density and taxonomic diversity (Chochong and Varatharajan, 2004). The species composition and mean percentage of their occurrence in Chochong and Varatharajan's (2004) study indicates the following sequence: *Urothrips tarai* (Stannard) - 20%, *Apelaunothrips madrasensis* (Ananthakrishnan) - 18%, *Adraneothrips okajimai* (Muraleedharan and Sen) - 12%, *Apelaunothrips consimilis* (Ananthakrishnan) and *Mecynothrips simplex* Bagnall - 10% each, *Elaphrothrips spiniceps* Bagnall, *Xylaplothrips debilis* Ananthakrishnan and Jagdish, and *Holurothrips manipurensis* Varatharajan and Chochong - 5% each, *Meiothrips menoni* Ananthakrishnan, *Bradythrips hesperus* Hood and Williams, and *Tylothrips indicus* Sen and Muraleedharan - 3% each, *Adraneothrips disjunctus* Ananthakrishnan, *Preeriella formosana* Okajima, and *Bactrothrips idolomorphus* Karny - 2% each (all Phlaeothripidae). Although the above information pertains to the *Quercus* forests of Manipur (24°44'N, 93°58'E), *U. tarai* (Phlaeothripidae) was first collected and described by Lewis Judson Stannard of the University of Illinois-Champaign, Illinois, in 1970, from the *Quercus* regions of the Tarai (29°22'N, 79°27'E). But the distribution of *U. tarai* from 24°44'N, 93°58'E to 28°2'N, 96°E of the north-eastern region of India confirms their occurrence throughout the *Quercus* belt of the Sub-Himalayan region. Similarly, species such as *T. indicus* and *H. manipurensis* are known only from Manipur, but the remaining 10 mycophagous Phlaeothripidae known earlier from the biodiversity-rich Western Ghats (Ananthakrishnan, 1973) are presently known to be occurring in north-eastern India as well. Further, the

minute leaf-litter inhabiting *P. formosana* (~950 µm) is a new record for India, which was previously known only from Taiwan (23°41'N, 120°57'E) (<http://anic.ento.csiro.au/thrips/resources/Taiwan.htm>). The presence of these 14 species especially in the *Quercus* belt reiterates that the thrips composition in the north-eastern India is a 'mixed bag' with species known previously from the Western Ghats, some from the Sub-Himalayan region, and some from the Indo-Myanmar bioregion (Varatharajan et al., 2010).

In bamboo forests

Nearly 125 native and 11 exotic species of bamboos (Poaceae) belonging to 23 genera are known from India, of which more than 50% occur in north-eastern India (fsi.nic.in/isfr2017/isfr-bambooresource). The overall bamboo-growing area in the Indian subcontinent has been estimated at *c.* 9.6 m ha. Thysanoptera are key inhabitants of bamboo litter (Ananthakrishnan, 1973; Varatharajan, 2005). The litter-inhabiting Thysanoptera of the Western Ghats and north-eastern India include five species of idolothripines and 10 of phlaeothripines. The species recorded thus far include *Acallurothrips amplus* (Faure), *Allothrips bicolor* Ananthakrishnan, *A. montanus* Ananthakrishnan, *Elaphrothrips curvipes* Priesner and *Nesothrips brevicollis* (Bagnall) (all Idolothripinae). On the other hand, members of the Phlaeothripinae include species such as *Ablemothrips maxillatus* Ananthakrishnan, *Adraneothrips bambusae* (Ananthakrishnan), *A. limpidus* Ananthakrishnan, *Apelaunothrips madrasensis* (Ananthakrishnan), *Karyothrips melaleucus* (Bagnall), *Margaritothrips flavus* Bhatti, *M. sumatrensis* Priesner, *Mystrothrips dammermani* (Priesner), *Ocythrips rarus* Ananthakrishnan, and *Stephanothrips occidentalis* Hood and Williams (Ananthakrishnan and Sen, 1980; Sen et al. 1988; Varatharajan, 2005).

In pine forests

A dozen mycophagous Phlaeothripidae occurs in association with the dry needle litter of *Pinus kesiya* Royle ex Gordon (Pinales: Pinaceae), of which *Apelaunothrips consimilis* (Ananthakrishnan), *Ecacanthothrips tibialis* (Ashmead), *Karyothrips melaleucus* (Bagnall), *Hoplandrothrips corticis* Ananthakrishnan, *Hoplothrips fungosus* Moulton, *Hoplothrips orientalis* (Ananthakrishnan), *Macrophthalmothrips splendidus* Ananthakrishnan, *Streptothrips orientalis* (Ananthakrishnan) (Phlaeothripinae) are more dominant than the spore-feeding thrips such as *Ethirothrips longisetis* (Ananthakrishnan and Jagdish), *Holothrips minor*

Table 1. List of Idolothripinae recorded in India

1. <i>Acallurothrips amplus</i> (Faure)	16. <i>Ethirothrips anacardii</i> (Ananth.)	31. <i>Elaphrothrips procer</i> (Schmutz)	44. <i>Meiothrips nepalensis</i> Kudo & Ananthakrishnan
2. <i>Aesthesiothrips jatrophae</i> Ananthakrishnan	17. <i>Ethirothrips beelsoni</i> (Moulton)	32. <i>Elaphrothrips spiniceps</i> Bagnall.	45. <i>Neosmerinthothrips fructuum</i> Schmutz
3. <i>Allothrips bicolor</i> Ananthakrishnan	18. <i>Ethirothrips brevisetosus</i> (Ananthakrishnan & Jagadish)	33. <i>Gastrothrips acuticornis</i> (Hood)	46. <i>Neosmerinthothrips inquilinus</i> Ananthakrishnan
4. <i>Allothrips indicus</i> Ananthakrishnan	19. <i>Ethirothrips brevis</i> (Bagnall)	34. <i>Gastrothrips falcatus</i> (Ananthakrishnan)	47. <i>Neosmerinthothrips robustus</i> (Ananthakrishnan)
5. <i>Allothrips montanus</i> Ananthakrishnan	20. <i>Ethirothrips indicus</i> (Bagnall)	35. <i>Gastrothrips turbinatus</i> (Ananthakrishnan)	48. <i>Nesidiothrips alius</i> (Ananthakrishnan)
6. <i>Allothrips watsoni</i> Hood	21. <i>Ethirothrips longisetis</i> (Ananthakrishnan & Jagadish)	36. <i>Holurothrips manipurensis</i> Varatharajan & Chochong	49. <i>Nesothrips brevicollis</i> (Bagnall)
7. <i>Bactrothrips idolomorphus</i> (Karny).	22. <i>Ethirothrips obscurus</i> (Schmutz)	37. <i>Ischyrothrips crassus</i> Schmutz	50. <i>Nesothrips lativentris</i> (Karny)
8. <i>Bactrothrips luteus</i> Ananthakrishnan	23. <i>Ethirothrips uredinis</i> (Ananthakrishnan & Jagadish)	38. <i>Loyolaia indica</i> Ananthakrishnan	51. <i>Ophthalmothrips breviceps</i> (Bagnall)
9. <i>Compsothrips congoensis</i> (Hood).	24. <i>Ethirothrips vitreipennis</i> (Priesner)	39. <i>Machatothrips corticosus</i> Ananthakrishnan	52. <i>Ophthalmothrips faurei</i> (Ananthakrishnan)
10. <i>Compsothrips ramamurthii</i> (Ananthakrishnan)	25. <i>Ethirothrips tirumalaiensis</i> (Ananthakrishnan)	40. <i>Machatothrips indicus</i> Ananthakrishnan and Jagadish	53. <i>Priesneriana kabandha</i> (Ramakrishna)
11. <i>Diaphorothrips unguipes</i> Karny	26. <i>Elaphrothrips curvipes</i> Priesner	41. <i>Machatothrips silvaticus</i> Ananthakrishnan	54. <i>Tiarothrips subramanii</i> (Ramakrishna)
12. <i>Dinothrips juglandis</i> Moulton	27. <i>Elaphrothrips denticollis</i> (Bagnall)	42. <i>Mecynothrips simplex</i> Bagnall	
13. <i>Dinothrips longicauda</i> (Ananthakrishnan).	30. <i>Elaphrothrips notabilis</i> Ananthakrishnan	43. <i>Meiothrips menoni</i> Ananthakrishnan	
14. <i>Dinothrips spinosus</i> (Schmutz)			
15. <i>Dinothrips sumatrensis</i> Bagnall			

(Hood), *Machatothrips indicus* Ananthakrishnan, and *Mecynothrips simplex* Bagnall (Idolothripinae). Among these, the bark-dwelling *Hoplandrothrips corticis* were apparently restricted to dry fallen bark pieces of *P. kesiya* that grow above 1500 masl, whereas the other species such as *H. fungosus* and *H. orientalis* inhabit the dry, fungus-infested needle litter. The density and diversity of Thysanoptera collected from the needle litter of *P. kesiya* was lesser than the other plantation sectors such as bamboo and oak (Table 3).

B. Thrips-plant-fungus association

Mycophagous Thysanoptera feed on fungal spores and hyphae and use dry leaves and twigs for egg laying and colony establishment. The mycophagous Phlaeothripidae mostly show a pattern in their colonization behaviour and feeding on fungal spores. For instance, the spore feeder *Tiarothrips subramanii* (Idolothripinae) occurs closely associated with three

species of *Anthostomella* (Ascomycota: Xylariales: Xylariaceae), that usually grows on dry fronds of *Borassus flabellifer* L. (Arecaceae). Similarly, *Mecynothrips hardyi* (Phlaeothripidae) feeds on the spores of *Dothiorella thripsita* Shivas and Tree (Ascomycota: Botryosphaerales: Botryosphaeriaceae) (Shivas et al., 2009). Species of *Pestalotia* (47.44%) (Ascomycota: Xylariales: Amphisphaeriaceae) have been found in the dissected gut of *B. idolomorphus* indicate the preference between *B. idolomorphus* and the species of *Pestalotia*. Further, *B. idolomorphus* also appear to prefer spores of the Coelomycetes, Ascomycetes, and Hyphomycetes, but not those of the Basidiomycetes (Ananthakrishnan and Dhileepan, 1984). Such a pattern is evident in spore- and mycetophagous species (Table 4), based on field observations and laboratory evaluation of spores of the gut contents. But, in Nature, a chance of more than one fungal taxon co-occurring at the same feeding site is highly likely.

Table 2. List of fungal-hyphae feeding Phlaeothripinae in India

1.	<i>Ablemothrips maxillatus</i> Ananthakrishnan	52.	<i>Hoplothrips orientalis</i> (Ananthakrishnan)
2.	<i>Adraneothrips bambusae</i> (Ananthakrishnan)	53.	<i>Hoplothrips transvaalensis</i> (Hood)
3.	<i>Adraneothrips disjunctus</i> Ananthakrishnan	54.	<i>Idiothrips bellus</i> Faure
4.	<i>Adraneothrips elegans</i> Ananthakrishnan	55.	<i>Karnyothrips melaleucus</i> (Bagnall)*
5.	<i>Adraneothrips infirmus</i> (Ananthakrishnan)	56.	<i>Karnyothrips mucidus</i> (Ananthakrishnan & Jagdish)*
6.	<i>Adraneothrips limpidus</i> (Ananthakrishnan)	57.	<i>Macrophthalthmothrips splendidus</i> Ananthakrishnan
7.	<i>Adraneothrips madrasensis</i> Ananthakrishnan	58.	<i>Malacothrips natalensis</i> (Trybom)
8.	<i>Adraneothrips nilgiriensis</i> (Ananthakrishnan)	59.	<i>Margaritothrips flavus</i> Bhatti
9.	<i>Adraneothrips okajimai</i> (Muraleedharan & Sen)	60.	<i>Margaritothrips longus</i> Bhatti
10.	<i>Adraneothrips pteris</i> (Ananthakrishnan)	61.	<i>Margaritothrips sumatrensis</i> Priesner
11.	<i>Adraneothrips stannardi</i> (Ananthakrishnan)	62.	<i>Mesandrothrips emineus</i> (Ananthakrishnan & Jagdish)**
12.	<i>Apelaunothrips bhowalii</i> (Ananthakrishnan)	63.	<i>Mesandrothrips pictipes</i> (Bagnall)**
13.	<i>Apelaunothrips consimilis</i> (Ananthakrishnan)	64.	<i>Mesandrothrips pusillus</i> (Ananthakrishnan & Jagdish)**
14.	<i>Apelaunothrips indicus</i> (Ananthakrishnan)	65.	<i>Mesandrothrips tener</i> (Ananthakrishnan & Jagdish)**
15.	<i>Apelaunothrips lucidus</i> (Ananthakrishnan)	66.	<i>Mystrithrips dammermani</i> (Priesner)
16.	<i>Apelaunothrips madrasensis</i> (Ananthakrishnan)	67.	<i>Neothrips lepidus</i> Ananthakrishnan
17.	<i>Apterygothrips fungosus</i> (Ananthakrishnan & Jagdish)*	68.	<i>Ocythrips rarus</i> Ananthakrishnan
18.	<i>Apterygothrips jogensis</i> (Ananthakrishnan & Jagdish)*	69.	<i>Oidanothrips enormis</i> (Ananthakrishnan)
19.	<i>Apterygothrips rubiginosus</i> (Ananthakrishnan & Jagdish)*	70.	<i>Oidanothrips megacephalus</i> (Ananthakrishnan)
20.	<i>Azaleothrips amabilis</i> Ananthakrishnan	71.	<i>Opidnothrips corticulus</i> Ananthakrishnan
21.	<i>Azaleothrips aspersus</i> Bhatti	72.	<i>Phiarothrips reperticus</i> Ananthakrishnan
22.	<i>Azaleothrips bhattii</i> Vijay Veer & Chauhan	73.	<i>Phlaeothrips nilgircus</i> Ananthakrishnan
23.	<i>Azaleothrips lineus</i> Bhatti	74.	<i>Phylladothrips karnyi</i> Priesner
24.	<i>Baenothrips asper</i> (Bournier)	75.	<i>Plectrothrips corticinus</i> Priesner
25.	<i>Baenothrips indicus</i> (Ananthakrishnan)	76.	<i>Plectrothrips eximius</i> Ananthakrishnan
26.	<i>Baenothrips minutus</i> (Ananthakrishnan)	77.	<i>Plectrothrips pallipes</i> Hood
27.	<i>Bradythrips hesperus</i> Hood & Williams	78.	<i>Preeriella formosana</i> Okajima
28.	<i>Bunothrips cruralis</i> Ananthakrishnan.	79.	<i>Priesneria insolitus</i> (Ananthakrishnan)
29.	<i>Ecacanthothrips tibialis</i> (Ashmead)	80.	<i>Psalidothrips ascitus</i> (Ananthakrishnan)
30.	<i>Glubothrips mucidus</i> Ananthakrishnan	81.	<i>Pygmaeothrips angusticeps</i> (Hood)
31.	<i>Habrothrips curiosus</i> Ananthakrishnan	82.	<i>Socothrips verrucosus</i> Ananthakrishnan
32.	<i>Holothrips andamanensis</i> (Sen)	83.	<i>Sophiothrips nigrus</i> Ananthakrishnan
33.	<i>Holothrips ananthakrishnani</i> Okajima	84.	<i>Sophiothrips typicus</i> (Ananthakrishnan)
34.	<i>Holothrips cracens</i> (Ananthakrishnan)	85.	<i>Stannardothrips longirostris</i> Ananthakrishnan
35.	<i>Holothrips fumidus</i> (Ananthakrishnan)	86.	<i>Stephanothrips adnatus</i> Ananthakrishnan
36.	<i>Holothrips indicus</i> (Ananthakrishnan)	87.	<i>Stephanothrips occidentalis</i> Hood & Williams
37.	<i>Holothrips minor</i> (Hood)	88.	<i>Strepterothrips orientalis</i> Ananthakrishnan
38.	<i>Holothrips mirandus</i> (Ananthakrishnan)	89.	<i>Symphyothrips aberrans</i> Ananthakrishnan
39.	<i>Holothrips nepalensis</i> (Pelikán)	90.	<i>Tamilthrips pini</i> (Ananthakrishnan)*
40.	<i>Holothrips quadrisetis</i> Okajima	91.	<i>Trichinothrips breviceps</i> (Bagnall)
41.	<i>Holothrips ruidus</i> (Ananthakrishnan)	92.	<i>Tylothrips indicus</i> Sen & Muraleedharan
42.	<i>Holothrips stannardi</i> (Ananthakrishnan)	93.	<i>Tylothrips samirseni</i> Varatharajan, Singh & Bala
43.	<i>Holothrips subtilis</i> (Ananthakrishnan)	94.	<i>Urothrips tarai</i> (Stannard)
44.	<i>Holothrips typicus</i> (Ananthakrishnan)	95.	<i>Veerabahuthrips bambusae</i> Ramakrishna
45.	<i>Hoplandrothrips corticis</i> Ananthakrishnan	96.	<i>Xylaplothrips debilis</i> Ananthakrishnan & Jagdish*
46.	<i>Hoplandrothrips flavipes</i> Bagnall	97.	<i>Xylaplothrips ligs</i> Ananthakrishnan & Jagdish*
47.	<i>Hoplandrothrips kudo</i> Muraleedharan	98.	<i>Xylaplothrips micans</i> Ananthakrishnan & Jagdish*
48.	<i>Hoplandrothrips nobilis</i> Priesner		
49.	<i>Hoplothrips dubius</i> (Bagnall)		
50.	<i>Hoplothrips fungosus</i> Moulton		
51.	<i>Hoplothrips nemorius</i> Ananthakrishnan		

*Species under Haplothripini and the rest come under *Phlaeothrips* lineage; ** *Mesandrothrips* (= *Xylaplothrips*)

Table 3. Thrips diversity and density in the leaf litters of natural and human-made forests of Manipur (24°44'N, 93°58'E)

Plantation type with altitude	Number of thrips species collected			Mean No.*
	Idolo- thripinae	Phlaeo- thripinae	Total	
Pine forest (1500-2100 m)	4	8	12	9.3 ^a
Bamboo forest (900 - 1600 m)	5	10	15	17.4 ^b
Oak forest (900-1300 m)	5	9	14	27.0 ^c
Polyculture forest (900-1800 m)	12	30	42	33.0 ^d

CD at p=0.01=4.6

Each value mean of seven replications*; Alphabets followed by each figure in the last vertical column different from each other at 1% level (ANOVA p=0.01); *No. collected using Tullgren funnel method

If this were proved right, then the spore-feeding thrips show no specificity to any particular fungal taxon.

Morpho-taxonomy distinguishes members of the Phlaeothripidae into spore-feeding Idolothripinae and the fungal-hyphae-feeding Phlaeothripinae, based on the width of maxillary stylets: 2-3 μm in the fungal-hypha feeders and 5-10 μm in spore feeders (Mound and Palmer, 1983). This classification gains in validity by the correlation of the width of maxillary stylets of

Mecynothrips hardyi (Priesner) (Phlaeothripidae) and fungal spore size of *Dothiorella thripsita* (Ascomycota: Botryosphaeriales: Botryosphaeriaceae). For example, the mean width of the maxillary stylet of *M. hardyi* (body length - 15 mm) is $13.9 \pm 0.2 \mu\text{m}$ and the mean width of the ingested spore of *D. thripsita* is $10.9 \pm 0.3 \mu\text{m}$. This observation along with the spores extracted from the gut confirmed the spore feeding habit of *M. hardyi* (Tree et al., 2010; Eow et al., 2011). This example serves well to illustrate mycophagous habit of the other Idolothripinae.

The mutualistic relations between insects and fungi in terms of dispersal, nutrition, mechanical protection and antimicrobial defence has been elaborately explained (Bidermann and Vega, 2019). Considering that in the context of Phlaeothripidae and fungi, the mutualism between them can be described in terms of dispersal and nutrition of the involved thrips. Nearly 2100 species of the Phlaeothripidae feed on fungal hyphae and spores (Dang et al., 2014; Eow et al., 2011). Some evidences also exist to indicate that the Thysanoptera act as mechanical vectors and disseminate fungal spores on plants (Ananthkrishnan, 1980; 1993). Mechanical dispersal of fungal spores is a strong possibility because of their numerous body setae that facilitate the attachment of fungal spores to their body surfaces similar to that of pollen grains transmitted mechanically by certain species of the Thysanoptera, such as *Thrips hawaiiensis* (Morgan), *Frankliniella schultzei*

Table 4. Thrips, plants and fungi

Thrips (Tubulifera: Idolothripinae)	Plant	Fungus
<i>Loyolaia indica</i> (Ananthkrishnan)	Dry leaves and leaf sheaths of <i>Cynodon dactylon</i> (L.) Pers. (Poaceae)	<i>Lojkania cynodontifolii</i> ⁺⁺ (Fenestellaceae)
<i>Tiarothrips subramanii</i> (Ramakrishna)	Dry leaves of <i>Borassus flabellifer</i> L. (Arecaceae)	<i>Anthostomella consanguinea</i> , <i>A. phoenicicola</i> , <i>A. sepebilibi</i> ⁺⁺ (Xylariaceae)
<i>Mecynothrips simplex</i> Bagnall	Dry leaves of <i>Areca catechu</i> L. (Arecaceae)	<i>Pestalotia</i> sp. ⁺⁺ (Amphisphaeriaceae)
<i>Mecynothrips hardyi</i> (Priesner)	<i>Acacia harpophylla</i> F. Muell. ex Benth. (Fabaceae)	<i>Dothiorella thripsita</i> ^{**} (Botryosphaeriaceae).
<i>Priesneriana kabandha</i> (Ramakrishna)	<i>Eucalyptus globulus</i> Labill. (Myrtaceae)	<i>Rhytidhysterium rufula</i> [*] (Patellariaceae) and <i>Cytospora</i> [*] (Phyllostictaceae)
<i>Elaphrothrips denticollis</i> (Bagnall)	<i>Tectona grandis</i> L. f. (Lamiaceae)	<i>Pestalotia</i> sp. [*] (Amphisphaeriaceae) <i>Phomopsis tectonae</i> (Diaporthaceae)
<i>Nesothrips indicus</i> Ananthkrishnan	Decaying scape of <i>Agave americana</i> L. (Asparagaceae)	<i>Anthostomella sphaeroidea</i> [*] (Xylariaceae)
<i>Dinothrips sumatrensis</i> Bagnall	Dry bark of <i>Piper nigrum</i> L. (Piperaceae)	<i>Lasiodiplodia theobromae</i> [*] (Botryosphaeriaceae)

Source: ⁺⁺Ananthkrishnan and James, 1983; ^{**}Tree et al. (2010) ^{*}Ananthkrishnan et al. (1984)

(Trybom), and *Microcephalothrips abdominalis* (Crawford) (Thripidae) (Varatharajan et al., 2016).

C. Developmental pattern

A comparison of breeding behaviours and developmental durations of different life stages of five species each of the mycetophagous and sporophagous thrips is available (Ananthakrishnan et al., 1984) (Table 5). The duration of development from egg to adult for the above 10 species was between 15 and 28 d, but the type of reproduction varied between the mycetophagous and sporophagous thrips (Ananthakrishnan et al., 1983; 1984). For instance, the mycetophagous Phlaeothripinae were oviparous and oviposited in greater numbers during moist months (July-December) when fresh fungi were abundantly available, whereas the sporophagous Idolothripinae reproduced by either oviparous or ovoviviparous or viviparous mode influenced by fungal density and environmental factors such as temperature, rainfall, and humidity. Field observations of the idolothripines such as *T. subramanii* and *B. idolomorphus* indicated viviparous mode of reproduction especially when the fungal density was less due to seasonal dryness. This has been confirmed by rearing *B. idolomorphus* at 30°C and 80% RH, wherein they reproduced either ovoviviparously or viviparously (Ananthakrishnan and Dhileepan, 1984). In *Ethirothrips*, another idolothripine, Ananthakrishnan et al. (1984) have observed that among the females of a natural population inhabiting the dry, fungus-infested leaves of a species of *Zizyphus* (Rhamnaceae), oviparous females formed 60% of that population, followed by viviparous and ovoviviparous females, 20% each. Similar to *B. idolomorphus*, *Bactrothrips* (= *Caudothrips*) *buffai* (Karny) (Idolothripinae) also reproduces ovoviviparously (Bournier, 1957). This behaviour has been interpreted as an adaptation wherein few larvae and eggs are laid in different batches

possibly to make use of available fungal material for the development of the juveniles (Bournier, 1957; 1966). But a comparison between oviparous and viviparous females in terms of eggs-larvae output revealed that the fecundity was less during the viviparous phase (7-11 larvae/ female) compared with the oviparous phase (25-33 eggs/ female) of *T. subramanii*. Whatever is the type of reproduction, from the perspective of their breeding behaviour, the mycophagous Phlaeothripidae inhabit fungus-rich parts of the litter and lay eggs in such a way that the fungal spores are accessible to the emergent larvae. In species such as *T. subramanii*, both adult males and females exhibit parental care by guarding rows of eggs until hatching (Ananthakrishnan et al., 1983; Ananthakrishnan and Suresh, 1983).

Populations of mycophagous Phlaeothripidae are regulated by the richness of fungi. The density of litter-inhabiting Phlaeothripidae is a useful, although indirect, measure to estimate fungal density: i.e., larger the thrips colony, greater is the diversity and density of the fungal flora, and the vice versa (Mound, 1974; 1977). In the mycophagous Phlaeothripidae, more apterous individuals than alates occur in the first 4-5 wk. With the dwindling of the availability of fungal spores, the situation reverses with more of alates, as shown in *Priesneriana kabandha* (Ramakrishna) (Idolothripinae) feeding on the spores of a species of *Cytospora* (Ascomycota: Sordariomycetes: Valsaceae). Thus, the abundance and depletion of fungal spores not only reveal the high and less abundance of the Phlaeothripidae, but also reflect the proportions of apterous and alate individuals within a colony (Ananthakrishnan et al., 1983). Between the two extremes, brachypterous individuals occur in many bark-dwelling Phlaeothripidae. Brachypterae (apterous) arise when the fresh fungi are available, whereas the macropterae (alates) occur more in numbers when fungal density declines (Hood, 1940; Bournier, 1961). Influence of food in alary polymorphism can be understood with *Hoplothrips fungi* Zetterstedt (Phlaeothripidae) feeding on a species of *Stereum* (Agaricomycetes: Russulales: Stereaceae) on dead wood. Apterous individuals were abundant as long as *Stereum* populations were high and when *Stereum* was replaced by a species of *Mucor* (Mucoromycota: Mucorales: Mucoraceae), a large number of winged males and females of *Hoplothrips fungi* arose (Mound, 2005). This example highlights not only the significance of fungi as food in the production and maintenance of apterous individuals, but indirectly highlights the specificity of *H. fungi* to a species of *Stereum*.

Table 5. Mycophagous Phlaeothripidae

Mycetophagous (Phlaeothripinae)	Sporophagous (Idolothripinae)
<i>Adraneothrips limpidus</i> Ananthakrishnan	<i>Bactrothrips idolomorphus</i> (Karny)
<i>Azaleothrips amabilis</i> Ananthakrishnan	<i>Elaphrothrips denticollis</i> (Bagnall)
<i>Ecacanthothrips tibialis</i> (Ashmead)	<i>Elaphrothrips procer</i> (Schmutz)
<i>Holothrips cracens</i> (Ananthakrishnan)	<i>Meiothrips menoni</i> Ananthakrishnan
<i>Hoplandrothrips flavipes</i> Bagnall	<i>Tiarothrips subramanii</i> (Ramakrishna)

Although alary polymorphism occurs among the mycophagous Phlaeothripidae (Figs. 1a, b), the development of apterous individuals seems to be an adaptation in the leaf-litter habitat for the following reasons:

- (a) Abundance of fungi enables the Phlaeothripidae to feed and breed. For instance, *Loyolaia indica* consumes 47,600 spores of *Fusarium oxysporum* f. sp. *cubeense* Smith, Snyder and Hansen (Sordariomycetes: Hypocreales: Nectriaceae) per adult and lays 4-6 eggs/female on clumps of *Cynodon dactylon* Pers. (Poaceae). The total development time is 13-18 d when reared at $30 \pm 2^\circ\text{C}$ and 72% RH (Ananthkrishnan et al., 1984).
- (b) In the fungus-feeding *Suocerathrips linguis* Mound and Marullo (1994) in Germany, occurrence of either partial or completely de-alate adult females was common in the colony and the process of wing shedding occurred immediately after mating (Moritz, 2002; Mound, 2005). This feature was explained as *S. linguis*'s commitment to establish the colony and parental care.
- (c) The mycophagous Phlaeothripidae utilize the conducive semi-permanent niche endowed with abundant fungi to their advantage and produce more apterous adults as in *H. fungi* on the fungus-infested dead wood (Mound, 2005).
- (d) The juvenile hormone (JH) titre will be higher in apterous insects than the alates in general (Roff, 1991). Higher JH titre would mean that the duration on insect development will be longer in apterous than alates. For example, the total duration of development in *Loyolaia indica* was reported to be 13-18 d when reared at $30 \pm 2^\circ\text{C}$ and 72% RH, but the study did not mention it for alate or apterous individuals (Ananthkrishnan et al., 1984). Although the bioassay on thrips's hormone levels has not been attempted so far, considering the role of JH, it is presumed that the extended duration of 5 d of development may refer to the duration of apterous and the short duration of 13 d may reflect the developmental period of winged individuals, since all the rearing conditions stated above were the same.
- (e) Periodical collection and enumeration of field populations of litter-inhabiting thrips also reflected the presence of alates, apterous, normal males and females, besides large and small males reflecting morphological variation (Ananthkrishnan, 1969; 1979; Mound and Palmer, 1983; Tree et al., 2010). As a result, in a colony, it is possible to observe




Fig. 1. a, b: Alate and Apterous *Holurothrips manipurensis*; c, d: small and large individuals of *Hoplothrips fungosus* (Scale bar = $500 \mu\text{m}$). Arrows indicate size variation of fore-femora. (Source: c, d. Ananthkrishnan, 1973).

a small proportion of large and small males, in addition to dominant population of normal-size males and females of the same species. Size variation among them is distinct as in *T. subramanii*, in which the length of third antennal segment of the small, normal, and large individuals of *T. subramanii* were 80, 160 and $240 \mu\text{m}$, respectively (Ananthkrishnan, 1973; 1979). Similar variations on the size of forefemora of *H. fungosus* (Fig. 1c and 1d) (Ananthkrishnan, 1973), two fold long preocular projection on the heads of large males of *T. subramanii*, three-fold increased width of the fore-femora of large males of *E. tibialis* (Ananthkrishnan, 2005) and thorn-like cheek setae of large males of *Mecynothrips kraussi* Palmer and Mound, (Mound, 2005) are a few notable examples of intraspecific variation in comparison with the respective species of their smaller males. Based on the available data pertaining to representative

examples of mycophagous Thysanoptera mentioned here, a general pattern of their development has been reconstructed (Fig. 2).

D. Phenotypic plasticity

Phenotypic plasticity is a kind of intraspecific variation among individuals belonging to same species, in which, some members of a population exhibit considerable morphological variations (Ananthkrishnan, 2005). Such a plasticity is common among the mycophagous Phlaeothripidae. As of the present, only observational evidences exist in support of the prevalence of phenotypic plasticity among the Thysanoptera, and experimental verifications are necessary to strengthen the phenotypic-plasticity theory (Mound, 2005). The literature abounds with many observational reports that many species of mycophagous Phlaeothripidae exhibit wide-ranging morphological traits, such as alary polymorphism, sexual dimorphism, and intra-specific variation of body organs. Ananthkrishnan (1979; 1984; 2005) and Mound (2005) have elaborately discussed this aspect based on morpho-taxonomy. The following remarks highlight some of the prominent features noticed among selected mycophagous Phlaeothripidae:

- Alary polymorphism refers to either apterous or




Fig 2. Developmental pattern in certain fungal spore feeding thrips

brachypterous or alate individuals. As evident in *H. manipurensis*, populations from *Quercus*-leaf litter in summer months (April-June) included more of alates, whereas in winter and spring months (November-March), *H. manipurensis* populations included more of apterous individuals in Manipur (24°44'N, 93°58'E). *Quercus griffithii* Hook f. et Thomson ex. Miq., and *Q. serrata* Murray (Fagaceae) being deciduous, accumulation of leaf litter is appreciably more in late autumn and winter compared with summer. Dense leaf litter coupled with low temperature, dampness, and short photoperiods facilitates the growth of fungus, which highly likely lead to the production of apterous individuals (Fig. 1a, b).

- In *Loyolaila indica*, reproductive diapause occurs in June-September on dry grass clusters, and the dispersal of alates take place later leading to the production of apterous individuals through oviparous mode of reproduction in Chennai (13°4'N, 80°14'E) climate. Variation in climatic conditions along with food scarcity results in the production of alates and with the arrival of north-east monsoon in October-November, fungal growth is activated, which in turn, facilitates the population build-up of apterous individuals (Ananthkrishnan, 1984).
- The large and small individuals within the male populations of *Ecacanthothrips tibialis*, *Elaphrothrips denticolis*, *M. simplex*, *T. subramanii*, are a few examples showing striking intraspecific structural variations within one population. Their allometric data show considerable differences between the large and small individuals. For instance, in *E. tibialis*, the width of fore-femora in large males is (1600-1800 μm) three times more than that of small male (400-600 μm). Similarly, the presence of fore-femora inner tooth (400-550 μm) can be observed in large males of *E. tibialis* and that will be almost invisible in small males of the same species (Ananthkrishnan, 1971). Although Ananthkrishnan (1971) has elaborately described this phenomenon in many a thrips species from a taxonomic perspective, Bernard Crespi (1986) conceptualized their functional dynamics and has explained such variations within the same colony. According to Crespi (1986), in a large congregation of fungus-feeding male *Hoplothrips pedicularius* Haliday (Phlaeothripinae) competitive infighting occurs resulting in the death of the weaker individuals. The successful males of *H.*

pedicularius feed voraciously and breed to protect their eggs and siblings (Crespi, 1988). Similar to other subsocial insects such as termites (Isoptera) and fig wasps (Hymenoptera) with traits to achieve the maximum level of inclusive fitness (Hamilton, 1978; 1979), spore-feeding thrips as well, display a range of features such as parental care, sexual dimorphism and other allometric variations among the individuals of a colony, possibly with the purpose to enhance their survival fitness.

E. Ecological implications

The taxonomic diversity of litter-inhabiting Thysanoptera is *c.* 2100 species feeding solely on fungal spores and hyphae (Dang et al., 2014; Eow et al., 2011). Not only their diversity, but their density are high, similar to populations of *Preeriella jacotia* Hartwig (Phlaeothripinae) with 2500 individuals collected from forest litter in South Africa (Hartwig, 1978). Such a diversity of species coupled with dense population possibly contributes to the overall ecological services to the concerned ecosystem. By virtue of occupying the litter habitat, the mycophagous Phlaeothripidae involve themselves in the decomposition and recycling of leaf litter, either directly or indirectly. Among the litter-inhabiting arthropods, the Phlaeothripidae occupy at least a third of the soil fauna. Although the Acarina are the most abundant (55.7%) soil arthropods, the Collembola and Thysanoptera standing second (14.7%) and third (6.5%), respectively, and the remainder including species of the Coleoptera, Hymenoptera, and the Lepidoptera (Wang and Tang, 2012). The role of litter-inhabiting thrips in forest floor is known to some extent (Mound, 1974; 1977; Ananthakrishnan, 1996), but, not many studies have been attempted to know the abundance and vertical distribution of insects in general and thrips in particular and such studies will certainly contribute substantially in understanding the role of litter-inhabiting Thysanoptera.

ACKNOWLEDGEMENTS

I am grateful to the Department of Science & Technology, New Delhi for financial support. This article is dedicated to N. Muraleedharan, former Director of the UPASI Tea Research Institute, Coimbatore & the Tocklai Tea Research Institute, Assam and Samir Sen, formerly Zoologist at the Zoological Survey of India, Calcutta for their contribution to the systematic studies of thrips of NE India. I am grateful to the Head, Department of Zoology and the coordinator, Centre of Advanced Study in Life Sciences, Manipur University

for the facilities. Thanks are due to LA Mound, CSIRO Entomology, Canberra, Australia for comments and suggestions.

REFERENCES

- Ananthakrishnan T N. 1969. Indian Thysanoptera. Council of Scientific & Industrial Research Zoological Monograph 1: 1-171.
- Ananthakrishnan T N. 1971. Trends in intraspecific sex-limited variation in some mycophagous Tubulifera (Thysanoptera). *Journal of the Bombay Natural History Society* 67: 481-501.
- Ananthakrishnan, T N. 1973. Mycophagous Tubulifera of India. Occasional Publications of the Entomology Research Institute, No. 2. 144 pp.
- Ananthakrishnan T N. 1979. Biosystematics of Thysanoptera. *Annual Review of Entomology* 24: 159-183.
- Ananthakrishnan T N. 1980. Thrips. Harris K F, Maramorosch K (eds.). *Vectors of plant pathogens*. Academic Press Inc., New York, 149-164 pp.
- Ananthakrishnan T N. 1984. *Bioecology of thrips*. Indira Publishing House, Michigan, 233 pp.
- Ananthakrishnan T N. 1993. Bionomics of thrips. *Annual Review of Entomology* 38: 71-92.
- Ananthakrishnan T N. 1996. *Forest litter insect communities: Biology and chemical ecology*, Oxford & IBH Publishing Co. Pvt. Ltd, New Delhi. 174 pp.
- Ananthakrishnan T N. 2005. Perspectives and dimensions of phenotypic plasticity in insects. Ananthakrishnan T N, Whitman D (eds.). *Insect phenotypic plasticity - diversity of responses*. Science Publishers, Inc., Enfield, 1-23 pp.
- Ananthakrishnan T N, Dhileepan, K. 1984. Thrips-fungus association with special reference to the sporophagous *Bactrothrips idolomorphus* (Karny) (Tubulifera: Thysanoptera). *Proceedings of the Indian Academy of Sciences (Animal Sciences)* 93: 243-249.
- Ananthakrishnan T N, James P W. 1983. Feeding preferences in some sporophagous Idolothropinae (Thysanoptera: Tubulifera). *Proceedings of the Indian Academy of Sciences (Animal Sciences)* B 49: 86-92.
- Ananthakrishnan T N, Sen S. 1980. *Taxonomy of Indian Thysanoptera*. Zoological Survey of India, Handbook Series No. 1, 234 pp.
- Ananthakrishnan T N, Suresh G. 1983. Patterns of fungal resource utilization and feeding range in some mycophagous Tubulifera (Insecta: Thysanoptera). *Proceedings. Indian Academy of Sciences (Animal Science)* 92: 285-291.
- Ananthakrishnan TN, Dhileepan K, Padmanabhan B. 1983. Reproductive strategies and behavioural attributes in some sporophagous Idolothropinae (Tubulifera: Thysanoptera). *Proceedings. Indian Academy of Sciences (Animal Sciences)* 92: 95-108.
- Ananthakrishnan T N, Padmanabhan B, Dhileepan K, Suresh G. 1984. Ecological interactions, species dynamics and reproductive biology of some mycophagous Thysanoptera. Occasional Publication No.5, Entomology Research Institute, Loyola College, Chennai. 1-37 pp.
- Bidermann P H W, Vega F E. 2020. Ecology and evolution of insect-fungus mutualisms. *Annual Review of Entomology* 65: 22.1-22.25.
- Bournier A. 1957. Un deuxième cas d'ovoviviparité chez les Thysanoptères, *Caudothrips buffai* Karny (Tubulifère, Megathripidae). *Comptes rendus de l'Académie des Sciences, Paris* 244: 506-508.
- Bournier A. 1961. Remarques au sujet du brachypterisme chez certaines espèces de thysanoptères. *Bulletin de la Société de Entomologie de France (NS)* 3: 173-179.

- Bournier A. 1966. L'embryogenèse de *Caudothrips buffai* Karny (Thysanoptera: Tubulifera). Annales de la Société entomologique de France (NS) 2: 415-435.
- Chochong V S, Varatharajan R. 2004. Faunistic diversity of thrips (Thysanoptera) inhabiting the oak leaf litters of plantation and natural habitats. Rajmohana K, Sudheer, K, Girish Kumar P, Santhosh S. (eds.). Perspectives on biosystematics and biodiversity, Prof T C Narendran commemorative volume, Systematic Entomology Research Scholars Association, Department of Zoology, University of Calicut, Calicut. 119-127 pp.
- Crespi B J. 1986. Territoriality and fighting in a colonial thrips, *Hoplothrips pedicularius*, and sexual dimorphism in Thysanoptera. Ecological Entomology 11: 119-130.
- Crespi B J. 1988. Adaptation, compromise and constraint: the development, morphometrics and behavioural basis of a fighter-flier polymorphism in male *Hoplothrips karnyi*. Behavioural Ecology and Sociobiology 23: 93-104.
- Dang L, Mound LA, Qiao G X. 2014. Conspectus of the Phlaeothripinae genera from China and Southeast Asia (Thysanoptera: Phlaeothripidae). Zootaxa 3807 (1): 1-82.
- Eow L X, Mound LA, Ng Y F. 2011. Genera of spore feeding Thysanoptera from Southeast Asia (Phlaeothripidae: Idolothripinae), with a species checklist from Peninsular Malaysia. Zootaxa 2928: 1-19.
- Hamilton W D. 1978. Evolution and diversity underbark, Mound LA, Waloff N (eds.). Diversity of insect faunas. Symposia of the Royal Entomological Society of London No 9, Oxford: Blackwell Scientific. 154-175 pp.
- Hamilton W D. 1979. Wingless and fighting males in fig wasps and other insects, Blum M S, Blum N A (eds.). Reproductive competition, mate choice and sexual selection in Insects, London, Academic Press. 167-220 pp.
- Hartwig E K. 1978. A new species of *Preeriella* (Thysanoptera: Phlaeothripidae) from South Africa with comments on Hydiotripini characters. Journal of Entomological Society of South Africa 41: 259-264.
- Hood J D. 1940. The cause and significance of macropterism and brachypterism in certain Thysanoptera, with description to a new Mexican species. Anales de la Escuela de Ciencias Biologicas 1: 497-505.
- Kirk W D J. 1996. Thrips: Naturalists' handbook. The Richmond Publishing Company 25: 1-70.
- Kobro S. 2010. Checklist of Nordic Thysanoptera. Norwegian Journal of Entomology 58: 21-26.
- Marren P, Mabey R. 2010. Bugs Britannica. Catalogue No.17419, Random House, London. 500 pp.
- Moritz G. 2002. The biology of thrips is not the biology of their adults: a developmental view. Marullo R, Mound L A (eds.). Thrips and tospoviruses, Proceedings. 7th International symposium on Thysanoptera. 259-267 pp.
- Morse J G, Hoddle M S. 2006. Invasion biology of thrips. Annual Review of Entomology 51: 67-89.
- Mound L A. 1974. Spore feeding thrips (Phlaeothripidae) from leaf litter and dead wood in Australia. Australian Journal of Zoology 27: 1-106.
- Mound L A. 1977. Species diversity and the systematics of some New World leaf litter Thysanoptera (Phlaeothripinae: Glyptothripini). Systematic Entomology 2: 225-244.
- Mound L A. 2002. Thysanoptera biodiversity in the Neotropics. Revista de Biologia Tropical 50: 477-484.
- Mound L A. 2005. Thysanoptera: diversity and interactions. Annual Review of Entomology 50: 247-269.
- Mound L A. 2013. Order Thysanoptera Haliday, 1836, in Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness (Addenda 2013). Zootaxa 3703 (1): 49-50
- Mound L A, Palmer J M. 1983. The generic and tribal classification of spore-feeding Thysanoptera (Phlaeothripidae: Idolothripinae). Bulletin of British Museum (Natural History), Entomology 46: 1-174.
- Mound L A, Tree D J. 2011. Australian spore-feeding Thysanoptera of the genus *Bactrothrips* (Phlaeothripidae: Idolothripinae). Zootaxa 3087: 56-65.
- Mound L A, Vesmanis A H. 2021. All genera of the world: Order Thysanoptera (Animalia: Arthropoda: Insecta), Megataxa 006 (1): 002-069.
- Mound L A, Dang, Li Hong, Tree D J. 2013. Genera of fungivorous Phlaeothripinae (Thysanoptera) from dead branches and leaf litter in Australia. Zootaxa 3681 (3): 201-224.
- Negi S S, Naithani H B. 1995. Oaks of India, Nepal and Bhutan. International Book Distributors, Dehradun. 266 pp.
- Okajima S. 1998. Minute leaf-litter thrips of the genus *Preeriella* (Thysanoptera, Phlaeothripidae) from Asia. Species Diversity 3: 301-316.
- Rachana R R, Varatharajan R. 2017. Checklist of terebrantian thrips (Insecta: Thysanoptera) recorded from India. Journal of Threatened Taxa 9: 9748-9755.
- Roff D A, 1991. Life history consequences of bio-energetic and biomechanical constraints on migration. American Zoologist 31: 205-215.
- Sen S, Pramanik N K, Sengupta C K. 1988. Thysanoptera fauna of North Eastern India, Records of the Zoological Survey of India, ZSI Occasional Paper No. 100. 123 pp.
- Shivas R G, Tree D J, Tan Y P, Ballard E L. 2009. *Dothiorella thripista*. Persoonia 22: 168-169.
- ThripsWiki, 2021(http://thrips.info/wiki/classification_overview) accessed on 17-5-2021.
- Tree D J, Mound L A, Walter G H. 2010. Fungal spore-feeding by adult and larval *Mecynothrips hardyi* (Priesner) (Thysanoptera: Phlaeothripidae: Idolothripinae). Journal of Natural History 44: 307-316.
- Tyagi K, Kumar V. (2016) Thrips (Insecta: Thysanoptera) of India - an updated checklist. Halteres 7: 64-98.
- Varatharajan R. 2005. Faunistic diversity of thrips (Thysanoptera) of north-eastern India. Silver Jubilee Publication, Manipur University. 74 pp.
- Varatharajan R, Tarunkumar O, James Keisa, T, Chochong V S, Taptamani H. 2010. Thysanoptera fauna of the oak fields of Manipur and Nagaland, north-eastern India. Entomon 35: 139-142.
- Varatharajan R, Shyam Maisnam, Chochong V S, Rachana R R. 2016. Pollination potential of thrips (Insecta: Thysanoptera)-an overview. Zoo's Print XXXI: 6-12.
- Wang J, Tang X L. 2012. Species diversity, species dynamics and vertical distribution of litter dwelling thrips in an urban forest remnant of South China. Journal of Insect Science 12: 1-12.

(Manuscript Received: July, 2021; Revised: July, 2021;

Accepted: July, 2021; Online Published: September, 2021)

Online published (Preview) in www.entosocindia.org Ref. No. e21200