



## PERSISTENCE AND DEGRADATION BEHAVIOUR OF DIMETHOATE IN GRAPES

**BANKA KANDA KISHORE REDDY, K BHUVANESWARI\*, M PARAMASIVAM, A SUGANTHI, P GEETHA<sup>1</sup>**

Department of Agricultural Entomology; <sup>1</sup>Centre for Post Harvest Technology,  
Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India

\*Email: bhuvanewari.k@tnau.ac.in (corresponding author)

### ABSTRACT

Through a field experiment, the dissipation pattern of dimethoate in grapes was evaluated during December 2020 - February 2021. Dimethoate 30%EC was sprayed twice at ten-days interval at recommended (X) dose (445 g a.i ha<sup>-1</sup>) and double the recommended (2x) dose (890 g a.i ha<sup>-1</sup>). The samples were collected from 0 (2hr), 1, 3, 5, 7, 10, 15 and 20 days or till degradation to below detectable level (BDL) and at harvest time. The residues were extracted by the modified QuEChERS method and analyzed by LCMS. The method performance was satisfactory in terms of SANTE guidelines and with good linearity ( $r^2 > 0.99$ ). The mean total dimethoate residue including omethoate was 1.047 and 2.168 mg kg<sup>-1</sup> at x and 2x doses, respectively with half-lives of 5.47 and 5.59 days. The calculated Risk Quotient (RQ) at both x and 2x dose with Good Agricultural Practices (GAP) indicated that for dimethoate these are not safe for human health due to the intake of residue.

**Key words:** Grapes, dimethoate, omethoate, method validation, QuChERS, dissipation, risk assessment, risk quotient, acceptable daily intake, maximum residue level

Organophosphorus (OP) pesticides account for 34% of global insecticide consumption and are the most extensively used broad spectrum insecticides. There are around 100 different OP pesticides that are used to control pests in horticultural crops (Jaipieam et al., 2009). Among the OP pesticides, dimethoate [O, O-dimethyl S-(2-(methylamino)-2-oxoethyl) phosphorodithioate] is a broad spectrum systemic insecticide largely used to manage pests in fruits and vegetables (Zheng and Sun, 2014). Dimethoate is transformed to omethoate in crops due to the oxidation process, which is more harmful than dimethoate itself (EFSA, 2013).

In India, dimethoate is registered for control of aphids, mealybugs, hoppers and stem borer in various fruit crops viz., mango, banana, citrus, apple, fig and apricot but not registered for use in grapes (CIBRC, 2021). However, a survey conducted in major grape growing districts of Tamil Nadu revealed that 93.33% of farmers are using dimethoate as a plant protection input to mitigate the pest problem in grapes (Jayabal et al., 2020). Dimethoate dissipation has been widely investigated in various fruit crops viz., mango (Bhattacharjee and Dikshit, 2016), pomegranate (Utture et al., 2012) and guava (Khan et al., 2009; Devi et al., 2016). It is one of the most widely used insecticides in Indian viticulture and globally for pest management

(Moyer and Neal, 2014; Patil et al., 2017; Preetha and Stanley, 2020).

In a desperate bid to save the crop, farmers often take intensive sprays of dimethoate at the berry initiation stage which is likely to leave toxic residues in/ on berries and may be unsafe to the consumers. The consumption of insecticide treated product become risky due to the residual persistence of the insecticide. So, it is necessary that insecticide should be active against pests while leaving only tolerable residue on food commodity. OPs are common pollutants that pose significant toxicological risks to soil, aquatic ecosystems, and human health (Ji et al., 2019). Therefore, it is important to understand how pesticide residues dissipate at various stages in order to ensure the supply of safe and fresh produce to the end users.

To our knowledge, there are no published reports in India on the dissipation of dimethoate and its metabolite omethoate in grapes. Therefore, a field study was conducted to monitor the degradation pattern of dimethoate in response to the above issue. Using Indian dietary data, the preharvest interval (PHI) and dietary risk assessment were calculated.

### MATERIALS AND METHODS

For the estimation of dimethoate and omethoate

residues, certified reference material (CRM) of dimethoate (99.5%), omethoate (96.8%), anhydrous sodium citrate dibasic sesquihydrate ( $\geq 99\%$  purity) (Sigma Aldrich, Bangalore, India); HPLC grade of ethyl acetate ( $\geq 99.9\%$ ), n-hexane ( $\geq 99.9\%$ ) (Sisco Research Laboratories, Mumbai, India); LCMS grade methanol ( $> 99.9\%$ ) (Avantor Sciences, PA, USA); Anhydrous sodium chloride ( $\geq 99\%$ ), anhydrous tri-sodium citrate dihydrate ( $\geq 99\%$ ) (Merck, Mumbai, India); anhydrous magnesium sulphate ( $\geq 99\%$ ) (Himedia Laboratories, Mumbai, India); primary secondary amine (PSA, 40  $\mu\text{m}$ ) and graphitised carbon black (GCB) (Agilent Technologies, USA); and formic acid ( $\geq 99\%$ ) (Fisher Scientific Limited, Czech Republic) were obtained. The commercial formulation of dimethoate 30% EC was purchased locally from a pesticide dealer in Coimbatore, Tamil Nadu, India.

Individual stock solutions of 400 mg/l dimethoate and omethoate were made in LCMS grade methanol by independently weighing 10.05 and 10.33 mg of the analytical standards into a calibrated glass A volumetric flask (25 ml). Dimethoate and omethoate secondary stock solutions (40 mg/l) were made separately from the stock solution in a 25 ml container by transferring 2.5 ml of each. A working standard mixture of 10 mg/l was prepared using secondary stock solution. The linearity and spiking standard solutions were obtained by serial dilution from the mixed standard solution in the range of 0.005 - 0.1 mg/l. All standard solutions were maintained at  $-20^\circ\text{C}$  in the deep freezer until use. Matrix matched standard solutions (0.01, 0.025, 0.05, 0.075, and 0.1 mg/l) were prepared with immature and mature grape matrix.

Dissipation study was carried out in a farmer's field in Theni district of Tamil Nadu, India ( $9^\circ\text{N}$ ,  $76^\circ\text{E}$  and 375 masl) following good agricultural practices. The plot size was 50m<sup>2</sup> area that had no application of dimethoate before taking up the study, with three replicated plots. The commercial product of dimethoate 30% EC was applied (Muscat Hamburg variety) at the berry initiation stage (45 days after flowering) at the recommended dose (445 g a.i ha<sup>-1</sup>) and double the recommended dose (890 g a.i ha<sup>-1</sup>). A high-volume sprayer with 500 l ha<sup>-1</sup> of spray fluid, was used and two foliar sprays were given at an interval of 10 days for dissipation study. During the field trial, maximum and minimum temperatures were recorded as 28.16 and 19.16 $^\circ\text{C}$ , respectively and relative humidity was 78.16%. No rainfall was received.

The grape berries for residue analysis (0.5 kg) were

randomly picked from vines from each replication at 0 (2hr), 1, 3, 5, 7, 10, 15, 20, 25, 30 and 35 days after second spraying and at the time of harvest. A high-volume blade homogeniser was used to homogenise samples using a Robot Coupe cutter mixer (Blixer 6 VVA, France) and stored at  $-20^\circ\text{C}$  for residue analysis,

A modified QuEChERS (Quick, Easy, Cheap, Effective, Rugged, and Safe) technique was adapted, validated and used for residue extraction (Anastassiades et al., 2003). 10 g of grape sample was added in a 50 ml centrifuge tube and vortexed for one min after adding 20 ml of ethyl acetate. After that, 1 g NaCl and 4 g anhydrous MgSO<sub>4</sub> were added, vortexed and centrifuged completely for 10 min at 6000 rpm. Following centrifugation, a 6 ml aliquot of the supernatant was transferred to a 15 ml centrifuge tube containing 100 mg PSA, 600 mg anhydrous MgSO<sub>4</sub>, and 10 mg GCB and vortexed for one min before centrifugation at 3000 rpm for ten min. In a low volume concentrator at  $40^\circ\text{C}$ , 2 ml of aliquot was evaporated to near dryness under a gentle stream of nitrogen gas. Finally, the residue was mixed in 1.0 ml methanol and filtered by using a 0.2  $\mu\text{m}$  membrane syringe filter and transferred into 1.0 ml autosampler vials for analysis of residues by LC-MS.

The detection, estimation and confirmation of dimethoate and omethoate were done in Shimadzu 2020 series LCMS equipped with a reverse phase C18 (Eclipse plus- Agilent) column (250 mm length x 4.6 mm id, 5  $\mu\text{m}$  particle size) at a column oven temperature of  $40^\circ\text{C}$ . The mobile phase used for the separation of target compounds were methanol and ultra-pure water with 0.05 % formic acid in the ratio of 50:50. The LCMS pump was run in binary mode at a pressure of 48 kgf/cm<sup>2</sup> to discharge the mobile phase at a constant isocratic flow rate of 0.5 ml/min. The chromatograms were analysed using Shimadzu lab solutions software (5.6) and further calculations were based on obtained peak areas in the chromatograms. Samples were ionized using positive electron spray ionization (ESI+) mode and 0.1  $\mu\text{A}$  of interface current,  $350^\circ\text{C}$  of heat block temperature,  $250^\circ\text{C}$  as desolvation line (DL) temperature, nebulizer gas (N<sub>2</sub>-99.99%) flow of 1.5 l/min, drying gas of 12 l/min and scan speed of 15000 sec were employed.

The method used to determine the residues of dimethoate and omethoate in grape matrices was validated as per SANTE guidelines (SANTE, 2019) and evaluated with parameters viz., linearity, limit of detection (LOD), limit of quantification (LOQ),

recovery, precision, repeatability and matrix effect. The Horwitz ratio (HorRat) is a measurement tool that reflects the acceptability of analytical processes in terms of interlaboratory precision (reproducibility). It is the ratio of the observed relative standard deviation among laboratories (RSD), to the corresponding predicted relative standard deviation (PRSD), calculated using the Horwitz equation  $PRSD = 2C^{-0.15}$ , where C is the concentration found or added, expressed as a mass fraction and PRSD was calculated for all the fortified concentrations of 0.01, 0.025, 0.05, 0.075 and 0.1ppm (Horwitz and Albert, 2006). The matrix effect (ME) was assessed using the formula mentioned by (Dong et al., 2018).

The amount of dimethoate residue (log value) over time (days after application) was subjected to weighted linear regression to calculate half-life and safe waiting period values of dimethoate (Hoskins, 1961; Handa et al., 1999). Half-life of dimethoate was determined as  $T_{1/2} = 0.693/k$ . The maximum residue limit (MRL) for dimethoate in grapes was 0.01 mg/kg as per the European pesticide (EU) database, 2020. The safe waiting period of dimethoate was calculated using the formula mentioned as  $PHI = [\log(A) - \log(MRL)]/K$ .

EDI was divided by the acceptable daily intake (ADI) to arrive the risk quotient (RQ), which was used to quantify the long-term risk of intakes in comparison to pesticide toxicological data (Dong et al., 2018). ADI value of dimethoate is 0.002 mg/kg/bw/day (EFSA, 2013). The average body weight of an Indian (male- 65kg and female- 55kg) adult (NIN, 2020) and recommended total fruit consumption is 150 g/day (NIN, 2020). If the RQ is <1, the risk of long-term human dietary consumption of dimethoate is acceptable; if the RQ is >1, the risk is unacceptable.

## RESULTS AND DISCUSSION

Instrument conditions were optimised using single quadrupole LC-MS to identify, confirm, and quantify dimethoate and its metabolite omethoate in grapes. By setting standard chromatographic conditions, dimethoate and omethoate were separated and eluted at retention times of 8.47 and 5.31 min, respectively (Fig. 1). Quantification of dimethoate and omethoate was carried out using selected ion monitoring (SIM) mode and sensitivity was increased by using positive (+) SIM mode with target m/z at 230 and 214 for dimethoate and omethoate, respectively.

The recovery %, RSD, and linearity of the analytical




Fig. 1. Standard chromatogram of dimethoate + omethoate (0.01ppm)

method used to estimate the residues of dimethoate and omethoate were calculated and validated. In both immature and mature grape matrices, the method's linearity was determined to be between 0.005 to 0.1 mg/l. For both solvent and matrix-matched calibration standards, the correlation coefficient of dimethoate and omethoate was >0.99 (Table 1). The LOD (at a signal-to-noise ratio greater than 3) and LOQ (at a signal-to-noise ratio greater than 10) were determined as 0.005 and 0.01 mg/kg, respectively. The proposed LOQ (0.01 mg/kg) of the method was in agreement with the MRL (0.01 mg/kg) value fixed by EU pesticide database. Dimethoate and omethoate recovery in grapes was between 70 and 120% confirming to SANTE (2019) guidelines. RSD (<20%), matrix effect (<20%), horwitz ratio (0.5-2.0) were in acceptable range validating the analytical method (Table 1).

Residues of dimethoate reached BDL on 30 and 35 days after spraying at recommended (X) and double the recommended dose (2X), respectively (Table 2). Omethoate residues reached BDL 7 days after spraying. At harvest time samples, neither dimethoate nor omethoate was detected (45 days after spraying). Total residue (dimethoate + omethoate) was used to determine the half-life values and confirmed as 5.47 and 5.59 days at x and 2x doses, respectively. Reports on the dissipation of dimethoate in fruit crops like mango, guava, citrus, pomegranate are documented and available but could not get for grapes. Pappas et

Table 1. Recovery, linearity parameters and matrix effect of dimethoate and omethoate in grape matrices

Pesticides	Calibration (matrix)	Calibration range (mg/ l)	Regression equation	Correlation coefficient (R <sup>2</sup> )	Matrix effect (%)
Dimethoate	Solvent	0.005-0.1	y = 2E+07x + 39129	0.999	-
	Immature grapes	0.005-0.1	y = 2E+07x + 15789	0.9981	4.09
	Mature grapes	0.005-0.1	y = 2E+07x - 42273	0.9962	1.68
Omethoate	Solvent	0.005-0.1	y = 2E+07x + 36193	0.9993	-
	Immature grapes	0.005-0.1	y = 3E+07x + 13261	0.9972	3.56
	Mature grapes	0.005-0.1	y = 2E+07x - 4107.6	0.9985	6.79

Recovery of dimethoate and omethoate in grape matrix at different spiking levels (n=7)							
Matrix	Spiked concentration (mg/ kg)	Dimethoate			Omethoate		
		Recovery (%)± SD	RSD (%)	HorRat	Recovery (%)± SD	RSD (%)	HorRat
Immature grape berries	0.010	98.79± 2.49	2.52	0.08	96.59± 2.28	2.36	0.07
	0.025	93.55± 1.92	2.06	0.07	92.16± 2.71	2.94	0.11
	0.050	102.31± 1.64	1.60	0.06	103.05± 2.34	2.27	0.09
	0.075	97.92± 2.01	2.05	0.09	95.06± 1.80	1.89	0.08
	0.100	95.44± 2.90	3.04	0.14	95.61± 2.20	2.30	0.10
Mature grape berries	0.010	95.68± 1.72	1.80	0.06	85.28± 3.09	3.62	0.11
	0.025	90.10± 1.50	1.66	0.06	98.58± 1.48	1.51	0.05
	0.050	85.04± 1.68	1.98	0.08	93.54± 1.96	1.02	0.04
	0.075	92.08± 2.61	2.83	0.12	97.82± 2.88	2.94	0.13
	0.100	95.28± 3.17	3.33	0.15	94.19± 1.91	2.02	0.09

SD- standard deviation; RSD- relative standard deviation, HorRat- Horwitz ratio

Table 2. Residues, dissipation and dietary risk assessment of dimethoate in immature grape berries at 445 g a.i ha<sup>-1</sup> (X) and 890 g a.i ha<sup>-1</sup> (2X)

Days after treatment	x dose					2x dose					
	Dime-toate residues* (mg/ kg)	Ome-toate residues* (mg/ kg)	Total residues (mg/ kg)	Dietary risk assessment Risk quotient (RQ)		Days after treatment	Dime-toate residues* (mg/ kg)	Ome-toate residues* (mg/ kg)	Total residues (mg/ kg)	Dietary risk assessment Risk quotient (RQ)	
				Adult male (65 kg)	Adult female (55 kg)					Adult male (65 kg)	Adult female (55 kg)
0 (2hrs)	1.012	0.035	1.047	1.20	1.43	0 (2hrs)	2.107	0.061	2.168	2.50	2.96
1	0.607	0.026	0.633	0.73	0.86	1	1.098	0.051	1.149	1.32	1.56
3	0.527	0.022	0.549	0.63	0.75	3	0.802	0.035	0.837	0.96	1.14
5	0.385	0.011	0.396	0.46	0.54	5	0.632	0.028	0.660	0.76	0.90
7	0.272	BDL	0.272	0.31	0.37	7	0.567	0.014	0.581	0.67	0.80
10	0.162	BDL	0.162	0.19	0.22	10	0.337	BDL	0.337	0.39	0.46
15	0.099	BDL	0.099	0.11	0.13	15	0.295	BDL	0.295	0.34	0.40
20	0.078	BDL	0.078	0.09	0.11	20	0.205	BDL	0.205	0.24	0.28
25	0.044	BDL	0.044	0.05	0.06	25	0.091	BDL	0.091	0.10	0.12
30	0.014	BDL	0.014	0.01	0.01	30	0.051	BDL	0.051	0.06	0.07
35	BDL	BDL	BDL	-	-	35	0.010	BDL	0.010	0.01	0.01
Kinetic equation	y = 0.055x + 2.8862					y = 0.0538x + 3.1818					
R <sup>2</sup> value	0.974					0.947					
Half-life	5.47 days					5.59 days					
PHI	36.72 days					43.42 days					

BDL-Below Detectable Level (&lt;0.01 mg/ kg); \*Mean of three replicates, RQ- Risk Quotient

al. (2003) reported dissipation as 16.7 and 30.1 days in lemons and mandarins, respectively, 5.75 - 7 days in pomegranate (Utture et al., 2012), two days in mango (Bhattacharya and Diskhit, 2016) and 2.8 - 8.15 days in guava (Khan et al., 2009; Devi et al., 2016). The variation in the degradation pattern of dimethoate in different crops is due to the association of pesticide chemistry, plant architecture and environmental conditions in that particular crop ecosystem.

Risk assessment is the process of identifying potential threats and related hazards to life and health due to long term human exposure to chemicals found in food. The implication of total dimethoate residues including omethoate in grapes was evaluated by calculating risk quotient. The RQ value was >1 at both doses on the day of application and first day after spraying which indicated that grapes harvested on particular sampling days were not safe according to the calculated risk quotient value (RQ>1) under Tamil Nadu agroclimatic conditions.

#### ACKNOWLEDGEMENTS

Authors acknowledge Centre for Plant Protection Studies (CPPS) and Department of Agricultural Entomology, TNAU, Coimbatore for providing facilities.

#### REFERENCES

Anastassiades M, Lehotay S J, Stajnbaher D, Schenck F J. 2003. Fast and easy multiresidue method employing acetonitrile extraction / partitioning and “dispersive solid-phase extraction” for the determination of pesticide residues in produce. *Journal of AOAC International* 86(2): 412-431.

Bhattacharjee A K, Dikshit A. 2016. Dissipation kinetics and risk assessment of thiamethoxam and dimethoate in mango. *Environmental Monitoring and Assessment* 188(3): 165-169

CIBRC. Central Insecticide Board and Registration Committee. 2020. Available from [https://www.cibrc.ac.in/major\\_uses\\_of\\_pesticides](https://www.cibrc.ac.in/major_uses_of_pesticides). Accessed on 23.11.2020.

Devi M, Duhan A, Kumari B, Yadav G S. 2016. Determination of dimethoate, lambda-cyhalothrin and malathion residues in guava fruits using GCMS-tandem mass spectrometry. *Indian Journal of Horticulture* 73(2): 197-201.

Dong M, Wen G, Tang H, Wang T, Zhao Z, Song W, Wang W, Zhao L. 2018. Dissipation and safety evaluation of novaluron, pyriproxyfen, thiacloprid and tolfenpyrad residues in the citrus-field ecosystem. *Food chemistry* 269: 136-141.

EU Database. 2020. [https://ec.europa.eu/food/plant/pesticides/eu\\_pesticides-database/mrls](https://ec.europa.eu/food/plant/pesticides/eu_pesticides-database/mrls). Accessed 09.07.2020.

EFSA. European Food Safety Authority. 2013. Conclusion on the peer review of the pesticide risk assessment of confirmatory data submitted for the active substance dimethoate. *EFSA Journal* 11: 32-33.

Handa S, Agnihotri, Kulshrestha G. 1999. Maximum residue limits of pesticides. Pesticide residues significance, management and analysis. Research Periodicals and Book Publishing House, Houston. pp. 9-21.

Horwitz W, Albert R. 2006. The Horwitz ratio (HorRat): A useful index of method performance with respect to precision. *Journal of AOAC International* 89: 1095.

Hoskins W M. 1961. Mathematical treatment of the rate of loss of residues. Food and Agriculture Organization, FAO Plant Protection Bulletin 9: 163-168.

Jaipieam S, Visuthismajarn P, Sutheravut P, Siriwong W, Thomsang S, Borjan M, Robson M. 2009. Organophosphate pesticide residues in drinking water from artesian wells and health risk assessment of agricultural communities, Thailand. *Human and Ecological Risk Assessment* 15: 1304-1316.

Jayabal T D, Bhuvaneswari K, Reddy B K K. 2020. Survey on pest status, pesticide usage pattern in major grape growing districts of Tamil Nadu and recovery studies of select insecticides, Proceedings. International seminar on transboundary pest management, TNAU, Coimbatore. pp 338-339.

Ji Y, Wang Y, Yao Y, Ren C, Lan Z, Fang X, Zhang K, Sun W, Alder A C, Sun H. 2019. Occurrence of organophosphate flame retardants in farmland soils from northern China: Primary source analysis and risk assessment. *Environmental Pollution* 247: 832-838.

Khan B A, Farid A, Asi M R, Shah H, Badshah A K. 2009. Determination of residues of trichlorfon and dimethoate on guava using HPLC. *Food Chemistry* 114(1): 286-288.

Moyer M, Neal O S. 2014. Pest management strategic plan for Washington State wine grape production. Washington State University, Prosser, WA.

NIN. National Institute of Nutrition. 2020. [https://www.nin.res.in/RDA\\_Full\\_Report\\_2020.html](https://www.nin.res.in/RDA_Full_Report_2020.html). Accessed 02.07.2021.

Pappas C J, Kyriakidis N V. 2003. A comparison of dimethoate degradation in lemons and mandarins on the trees with two GC systems. *Food Chemistry* 80: 23-28

Patil N R, Nadaf A M, Patil D R. 2017. Field efficacy of newer insecticides and neem products against *Scirtothrips dorsalis* on grapes, *Vitis vinifera* L. (cv. Thompson Seedless). *International Journal of Current Microbiology and Applied Sciences* 6(9): 3287-3294.

Preetha G, Stanley J. 2020. Management of rose flower thrips, *Frankliniella schultzei* with insecticides. *Pesticide Research Journal* 32(1): 49-55.

SANTE. 2019. [http://ec.europa.eu/food/files/plant/docs/pesticides/mrl/guid\\_lines\\_wrkdoc.pdf](http://ec.europa.eu/food/files/plant/docs/pesticides/mrl/guid_lines_wrkdoc.pdf). Accessed 12.06.2020

Utture S C, Banerjee K, Kolekar S S, Dasgupta S, Oulkar D P, Patil S H, Wagh S S, Adsule P G, Anuse M A. 2012. Food safety evaluation of buprofezin, dimethoate and imidacloprid residues in pomegranate. *Food Chemistry* 131(3):787-795.

Zheng G L, Sun J L. 2014. Modern pesticide application technology series: insecticide. Chemical Industry Press, Beijing. 435 pp.